



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

eIF2B ϵ (m): 293T Lysate: sc-119973

BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. The eukaryotic initiation complex eIF2B exists as a five subunit complex composed of eIF2B α , eIF2B β , eIF2B γ , eIF2B δ and eIF2B ϵ . The eIF2B complex catalyzes the exchange of GDP for GTP on the eIF2 complex, following the interaction of eIF2/GTP with the 40S ribosomal subunit. Guanine nucleotide exchange factor (GEF) activity has been exhibited by the eIF2B ϵ subunit alone, but was greater in the presence of all five eIF2B subunits. Phosphorylation of eIF2 inhibits GEF activity of eIF2B, an inhibition that requires the eIF2B α subunit.

REFERENCES

1. Henderson, R.A., Krissansen, G.W., Yong, R.Y., Leung, E., Watson, J.D. and Dholakia, J.N. 1994. The δ -subunit of murine guanine nucleotide exchange factor eIF2B. Characterization of cDNAs predicts isoforms differing at the amino-terminal end. *J. Biol. Chem.* 269: 30517-30523.
2. Flowers, K.M., Mellor, H., Kimball, S.R. and Jefferson, L.S. 1995. Structure and sequence of the gene encoding the α -subunit of rat translation initiation factor-2B. *Biochim. Biophys. Acta* 1264: 163-167.
3. Price, N.T., Kimball, S.R., Jefferson, L.S. and Proud, C.G. 1996. Cloning of cDNA for the γ -subunit of mammalian translation initiation factor 2B, the guanine nucleotide-exchange factor for eukaryotic initiation factor 2. *Biochem. J.* 318: 631-636.
4. Price, N.T., Mellor, H., Craddock, B.L., Flowers, K.M., Kimball, S.R., Wilmer, T., Jefferson, L.S. and Proud, C.G. 1996. eIF2B, the guanine nucleotide-exchange factor for eukaryotic initiation factor 2. Sequence conservation between the α , β and δ subunits of eIF2B from mammals and yeast. *Biochem. J.* 318: 637-643.
5. Asuru, A.I., Mellor, H., Thomas, N.S., Yu, L., Chen, J.J., Crosby, J.S., Hartson, S.D., Kimball, S.R., Jefferson, L.S. and Matts, R.L. 1996. Cloning and characterization of cDNAs encoding the ϵ -subunit of eukaryotic initiation factor-2B from rabbit and human. *Biochim. Biophys. Acta* 1307: 309-317.
6. Webb, B.L. and Proud, C.G. 1997. Eukaryotic initiation factor 2B (eIF2B). *Int. J. Biochem. Cell Biol.* 29: 1127-1131.
7. Fabian, J.R., Kimball, S.R., Heinzinger, N.K. and Jefferson, L.S. 1997. Subunit assembly and guanine nucleotide exchange activity of eukaryotic initiation factor-2B expressed in Sf9 cells. *J. Biol. Chem.* 272: 12359-12365.

CHROMOSOMAL LOCATION

Genetic locus: Eif2b5 (mouse) mapping to 16 A3.

PRODUCT

eIF2B ϵ (m): 293T Lysate represents a lysate of mouse eIF2B ϵ transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

eIF2B ϵ (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive eIF2B ϵ antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.