



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



FTCD (m2): 293T Lysate: sc-120329

BACKGROUND

58K protein antibodies are excellent for use as markers for the Golgi complex. The 58K protein has been identified as being FTCD, a bifunctional enzyme that channels one-carbon units from formiminoglutamate, a metabolite of the histidine degradation pathway, to the folate pool. Defects in FTCD are the cause of glutamate formiminotransferase deficiency [also known as formiminoglutamyluria (FIGLU-uria)], an autosomal recessive disorder. Features of a severe phenotype include elevated levels of formiminoglutamate (FIGLU) in the urine in response to histidine administration, megaloblastic anemia and mental retardation. Features of a mild phenotype include high urinary excretion of FIGLU in the absence of histidine administration, mild developmental delay and no hematological abnormalities.

REFERENCES

- Hennig, D., Scales, S.J., Moreau, A., Murley, L.L., De Mey, J. and Kreis, T.E. 1998. A formiminotransferase cyclodeaminase isoform is localized to the Golgi complex and can mediate interaction of *trans*-Golgi network-derived vesicles with microtubules. *J. Biol. Chem.* 273: 19602-19611.
- Bashour, A.M. and Bloom, G.S. 1998. 58K, a microtubule-binding Golgi protein, is a formiminotransferase cyclodeaminase. *J. Biol. Chem.* 273: 19612-19617.
- Gao, Y.S., Alvarez, C., Nelson, D.S. and Sztul, E. 1999. Molecular cloning, characterization, and dynamics of rat formiminotransferase cyclodeaminase, a Golgi-associated 58 kDa protein. *J. Biol. Chem.* 273: 33825-33834.
- Gao, Y.S., Vrielink, A., MacKenzie, R. and Sztul, E. 2002. A novel type of regulation of the vimentin intermediate filament cytoskeleton by a Golgi protein. *Eur. J. Cell Biol.* 81: 391-401.
- Renous, R., Lapierre, P., Djilali-Saiah, I., Vitozzi, S. and Alvarez, F. 2003. Characterization of the antigenicity of the formiminotransferase-cyclodeaminase in type 2 autoimmune hepatitis. *Exp. Cell Res.* 292: 332-341.
- Mao, Y., Vyas, N.K., Vyas, M.N., Chen, D.H., Ludtke, S.J., Chiu, W. and Quiocho, F.A. 2004. Structure of the bifunctional and Golgi-associated formiminotransferase octamer. *EMBO J.* 23: 2963-2971.
- Hagiwara, H., Tajika, Y., Matsuzaki, T., Suzuki, T., Aoki, T. and Takata, K. 2006. Localization of Golgi 58K protein (formiminotransferase cyclodeaminase) to the centrosome. *Histochem. Cell Biol.* 126: 251-259.

CHROMOSOMAL LOCATION

Genetic locus: Ftc (mouse) mapping to 10 C1.

PRODUCT

FTCD (m2): 293T Lysate represents a lysate of mouse FTCD transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

FTCD (m2): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive FTCD antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


FTCD (G-3): sc-271788 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse FTCD expression in FTCD transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG_κ BP-HRP: sc-516102 or m-IgG_κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



FTCD (m2): sc-120329. Western blot analysis of FTCD expression in non-transfected: sc-117752 (**A**) and mouse FTCD transfected: sc-120329 (**B**) 293T whole cell lysates.



FTCD (G-3): sc-271788. Western blot analysis of FTCD expression in non-transfected: sc-117752 (**A**) and mouse FTCD transfected: sc-120329 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.