



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



GCM1 (m): 293T Lysate: sc-120453

BACKGROUND

GCM1 (glial cells missing homolog 1), also known as GCMA or hGCMA, is a 436 amino acid human homolog of the *Drosophila* glial cells missing protein (gcm). Localized to the nucleus and expressed specifically in placenta, GCM1 functions as a transcription factor that binds the novel sequence (A/G)CCCGCAT and, through this binding, regulates placental development. Additionally, GCM1 is thought to regulate syncytin SU-mediated trophoblastic fusion, an event that produces syncytiotrophoblast structures which, in turn, function as the outermost covering of the placental villi. GCM1 contains one N-terminal GCM (glial cell missing) DNA-binding domain, a conserved 150 amino acid residue that conveys DNA-binding activity for a variety of transcription factors involved in developmental processes.

REFERENCES

1. Akiyama, Y., Hosoya, T., Poole, A.M. and Hotta, Y. 1996. The gcm-motif: a novel DNA-binding motif conserved in *Drosophila* and mammals. Proc. Natl. Acad. Sci. USA 93: 14912-14916.
2. Yamada, K., Ogawa, H., Honda, S., Harada, N. and Okazaki, T. 1999. A GCM motif protein is involved in placenta-specific expression of human aromatase gene. J. Biol. Chem. 274: 32279-32286.
3. Yamada, K., Ogawa, H., Tamiya, G., Ikeno, M., Morita, M., Asakawa, S., Shimizu, N. and Okazaki, T. 2000. Genomic organization, chromosomal localization, and the complete 22 kb DNA sequence of the human GCMA/GCM1, a placenta-specific transcription factor gene. Biochem. Biophys. Res. Commun. 278: 134-139.
4. Yu, C., Shen, K., Lin, M., Chen, P., Lin, C., Chang, G.D. and Chen, H. 2002. GCMA regulates the syncytin-mediated trophoblastic fusion. J. Biol. Chem. 277: 50062-50068.
5. Baczyk, D., Satkunarathnam, A., Nait-Oumesmar, B., Huppertz, B., Cross, J.C. and Kingdom, J.C. 2004. Complex patterns of GCM1 mRNA and protein in villous and extravillous trophoblast cells of the human placenta. Placenta 25: 553-559.
6. Chang, C.W., Chuang, H.C., Yu, C., Yao, T.P. and Chen, H. 2005. Stimulation of GCMA transcriptional activity by cyclic AMP/protein kinase A signaling is attributed to CBP-mediated acetylation of GCMA. Mol. Cell. Biol. 25: 8401-8414.
7. Knerr, I., Schubert, S.W., Wich, C., Amann, K., Aigner, T., Vogler, T., Jung, R., Dötsch, J., Rascher, W. and Hashemolhosseini, S. 2005. Stimulation of GCMA and syncytin via cAMP mediated PKA signaling in human trophoblastic cells under normoxic and hypoxic conditions. FEBS Lett. 579: 3991-3998.
8. Chuang, H.C., Chang, C.W., Chang, G.D., Yao, T.P. and Chen, H. 2006. Histone deacetylase 3 binds to and regulates the GCMA transcription factor. Nucleic Acids Res. 34: 1459-1469.
9. Chou, C.C., Chang, C., Liu, J.H., Chen, L.F., Hsiao, C.D. and Chen, H. 2007. Small ubiquitin-like modifier modification regulates the DNA binding activity of glial cell missing *Drosophila* homolog a. J. Biol. Chem. 282: 27239-27249.

CHROMOSOMAL LOCATION

Genetic locus: Gcm1 (mouse) mapping to 9 E1.

PRODUCT

GCM1 (m): 293T Lysate represents a lysate of mouse GCM1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

GCM1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive GCM1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.