



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# MetAP-1 (m): 293T Lysate: sc-121609

## BACKGROUND

Methionine aminopeptidases (MetAP), also designated peptidase M proteins, are members of the M24 family of proteins. MetAP proteins remove the amino-terminal methionine residue from nascent polypeptides. MetAP-1 is a 394 amino acid protein that is expressed at low levels in all tissues, but is highly expressed in skeletal muscles. The active site of MetAP-1 contains two adjacent divalent metal ions connected by a water molecule or hydroxide ion. The control of cell proliferation in mammalian cells is directly linked and strictly dependent on the evolutionarily highly conserved mechanism that MetAP-1 employs. Eukaryotes contain both MetAP-1 and MetAP-2, whereas prokaryotes possess only the MetAP-1 enzyme. Pyridine-2-carboxylic acid thiazol-2-ylamide (PCAT) forms a scaffold that inhibits the action of MetAP-1, while 1,2,4-triazol is a non-peptide inhibitor of MetAP-1 binding to the active site with the N1 and N2 atoms of the triazole moiety complexing two divalent ions.

## REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 610151. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Oefner, C., et al. 2003. The 1.15Å crystal structure of the *Staphylococcus aureus* methionyl-aminopeptidase and complexes with triazole based inhibitors. *J. Mol. Biol.* 332: 13-21.
3. Brdlik, C.M. and Crews, C.M. 2004. A single amino acid residue defines the difference in ovalicin sensitivity between type I and II methionine aminopeptidases. *J. Biol. Chem.* 279: 9475-9480.
4. Swierczek, K., et al. 2005. Molecular discrimination of type-I over type-II methionyl aminopeptidases. *Biochemistry* 44: 12049-12056.
5. Bernier, S.G., et al. 2005. Methionine aminopeptidases type I and type II are essential to control cell proliferation. *J. Cell. Biochem.* 95: 1191-1203.
6. Luo, Q.L., et al. 2005. Inhibitors of type I MetAPs containing pyridine-2-carboxylic acid thiazol-2-ylamide. Part 1: SAR studies on the determination of the key scaffold. *Bioorg. Med. Chem. Lett.* 15: 635-638.

## CHROMOSOMAL LOCATION

Genetic locus: Metap1 (mouse) mapping to 3 G3.

## PRODUCT

MetAP-1 (m): 293T Lysate represents a lysate of mouse MetAP-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

MetAP-1 (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive MetAP-1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

MetAP-1 (A-2): sc-514653 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse MetAP-1 expression in MetAP-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



MetAP-1 (A-2): sc-514653. Western blot analysis of MetAP-1 expression in non-transfected: sc-117752 (A) and mouse MetAP-1 transfected: sc-121609 (B) 293T whole cell lysates.



MetAP-1 (C-10): sc-376779. Western blot analysis of MetAP-1 expression in non-transfected: sc-117752 (A) and mouse MetAP-1 transfected: sc-121609 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.