



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



pki α (m): 293T Lysate: sc-122609

BACKGROUND

The second messenger cyclic AMP (cAMP) mediates a diverse array of cellular responses such as proliferation, ion transport, regulation of metabolism and gene transcription by activating the cAMP-dependent protein kinase (cAPK or PKA). Activation of PKA occurs when cAMP binds to the two regulatory subunits of tetrameric PKA, resulting in the release of two active catalytic subunits. Two forms of a specific PKA inhibitor molecule, designated pki α and pki β , have been described and are suggested to regulate PKA activity in different portions of the central nervous system. For instance, pki α is expressed abundantly in the adult mouse brain, particularly in the cerebellum, hypothalamus, hippocampus and cortex. In contrast, pki β is present at a much lower level in most brain regions. pki β is found in significant amounts only in the cerebellum and in a few distinct nuclei within the pons, medulla and hypothalamus.

REFERENCES

1. Beavo, J.A., Bechtel, P.J. and Krebs, E.G. 1974. Activation of protein kinase by physiological concentrations of cyclic AMP. Proc. Natl. Acad. Sci. USA 71: 3580-3583.
2. Krebs, E.G. and Beavo, J.A. 1980. Phosphorylation and dephosphorylation of enzymes. Annu. Rev. Biochem. 48: 923-959.
3. Maldonado, F. and Hanks, S.K. 1988. cAMP-dependent protein kinase, α -catalytic subunit. Nucleic. Acids Res. 16: 8189-8190.
4. Beebe, S.J., Oyen, O., Sandberg, M., Froysa, A., Hansson, V. and Jahnse, T. 1990. cAMP-dependent protein kinase, β -catalytic subunit. Mol. Endoc. 4: 465-475.
5. Meinkoth, J.L., Alberts, A.S., Went, W., Fantozzi, D., Taylor, S.S., Hagiwara, M., Montminy, M. and Feramisco, J.R. 1993. Signal transduction through the cAMP-dependent protein kinase. Mol. Cell. Biochem. 127-128: 179-186.
6. Marchetto, G.S. and Henry, H.L. 1995. Cloning and sequencing of the cDNA encoding the avian kidney cAMP-dependent protein kinase inhibitor protein. Gene 158: 303-304.
7. Seasholtz, A.F., Gamm, D.M., Ballesteros, R.P., Scarpetta, M.A. and Uhler, M.D. 1995. Differential expression of mRNAs for protein kinase inhibitor isoforms in mouse brain. Proc. Natl. Acad. Sci. USA 92: 1734-1738.

CHROMOSOMAL LOCATION

Genetic locus: Pkia (mouse) mapping to 3 A1.

PRODUCT

pki α (m): 293T Lysate represents a lysate of mouse pki α transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

pki α (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive pki α antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.