



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



A-Raf (h): 293T Lysate: sc-159320



BACKGROUND

Several serine/threonine protein kinases have been implicated as intermediates in signal transduction pathways. These include ERK/MAP kinases, ribosomal S6 kinase (Rsk) and Raf-1. Raf-1 is a cytoplasmic protein with intrinsic serine/threonine activity. It is broadly expressed in nearly all cell lines tested to date and is the cellular homolog of v-Raf, the product of the transforming gene of the 3,611 strain of murine sarcoma virus. The unregulated kinase activity of the v-Raf protein has been associated with transformation and mitogenesis while the activity of Raf-1 is normally suppressed by a regulatory N-terminal domain. A-Raf, a second member of the Raf gene family of serine/threonine protein kinases, exhibits substantial homology to Raf-1 within the kinase domain of the two molecules, but less homology elsewhere. Expression of A-Raf is found at highest levels in urogenital tissues and kidney and at lowest level in brain tissue.

REFERENCES

1. Rapp, U.R., et al. 1983. Structure and biological activation of v-Raf, a unique oncogene transduced by a retrovirus. Proc. Natl. Acad. Sci. USA 80: 4218-4222.
2. Huleihel, M., et al. 1986. Characterization of murine A-Raf, a new oncogene related to the v-Raf oncogene. Mol. Cell. Biol. 6: 2655-2662.
3. Sariban, E., et al. 1987. Expression of the c-Raf proto-oncogene in human hematopoietic cells and cell lines. Blood 69: 1437-1440.
4. Ray, L.B., et al. 1988. Insulin-stimulated microtubule-associated protein kinase is phosphorylated on tyrosine and threonine *in vivo*. Proc. Natl. Acad. Sci. USA 85: 3753-3757.
5. Morrison, D.K., et al. 1988. Signal transduction from membrane to cytoplasm: growth factors and membrane-bound oncogene products increase Raf-1 phosphorylation and associated protein kinase activity. Proc. Natl. Acad. Sci. USA 85: 8855-8859.
6. Pelech, S.L., et al. 1990. Protein kinase cascades in meiotic and mitotic cell cycle control. Biochem. Cell Biol. 68: 1297-1330.
7. Heidecker, G., et al. 1990. Mutational activation of c-Raf-1 and definition of the minimal transforming sequence. Mol. Cell. Biol. 10: 2503-2512.
8. Turner, B., et al. 1991. Interleukin-2 induces tyrosine phosphorylation and activation of p72-74 Raf-1 kinase in a T cell line. Proc. Natl. Acad. Sci. USA 88: 1227-1231.

CHROMOSOMAL LOCATION

Genetic locus: ARAF (human) mapping to Xp11.23.

PRODUCT

A-Raf (h): 293T Lysate represents a lysate of human A-Raf transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

A-Raf (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive A-Raf antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.