



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



M-ficolin (h2): 293T Lysate: sc-159506

BACKGROUND

M-ficolin, also known as ficolin-A, ficolin-1, ficolin- α , FCN1, FCNM or collagen/Fibrinogen domain containing protein 1, is a 326 amino acid member of the ficolin lectin family. Characteristic of ficolin family proteins, M-ficolin contains a short N-terminus, one collagen-like domain and one Fibrinogen-like domain at its C-terminus. M-ficolin is a secreted innate immunity pattern recognition molecule predominantly expressed in peripheral blood leukocytes. It is believed to function as a plasma protein and is known to interact with elastin, carbohydrates and corticosteroids. M-ficolin exists as a homopolymer through intermolecular disulfide bonding and is involved in activating lectin activity via association with MASP and specific carbohydrate ligands.

REFERENCES

1. Ohashi, T. and Erickson, H.P. 1997. Two oligomeric forms of plasma ficolin have differential lectin activity. *J. Biol. Chem.* 272: 14220-14226.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 601252. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Hummelshøj, T., Munthe-Fog, L., Madsen, H.O., Fujita, T., Matsushita, M. and Garred, P. 2005. Polymorphisms in the FCN2 gene determine serum variation and function of Ficolin-2. *Hum. Mol. Genet.* 14: 1651-1658.
4. Frederiksen, P.D., Thiel, S., Larsen, C.B. and Jensenius, J.C. 2005. M-ficolin, an innate immune defence molecule, binds patterns of acetyl groups and activates complement. *Scand. J. Immunol.* 62: 462-473.
5. Liu, Y., Endo, Y., Iwaki, D., Nakata, M., Matsushita, M., Wada, I., Inoue, K., Munakata, M. and Fujita, T. 2005. Human M-ficolin is a secretory protein that activates the lectin complement pathway. *J. Immunol.* 175: 3150-3156.
6. Thiel, S. 2007. Complement activating soluble pattern recognition molecules with collagen-like regions, mannan-binding lectin, ficolins and associated proteins. *Mol. Immunol.* 44: 3875-3888.
7. Hummelshøj, T., Fog, L.M., Madsen, H.O., Sim, R.B. and Garred, P. 2007. Comparative study of the human ficolins reveals unique features of Ficolin-3 (Hakata antigen). *Mol. Immunol.* 45: 1623-1632.
8. Garlatti, V., Martin, L., Gout, E., Reiser, J.B., Fujita, T., Arlaud, G.J., Thielen, N.M. and Gaboriaud, C. 2007. Structural basis for innate immune sensing by M-ficolin and its control by a pH-dependent conformational switch. *J. Biol. Chem.* 282: 35814-35820.
9. Frankenberger, M., Schwaeble, W. and Ziegler-Heitbrock, L. 2008. Expression of M-ficolin in human monocytes and macrophages. *Mol. Immunol.* 45: 1424-1430.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: FCN1 (human) mapping to 9q34.3.

PRODUCT

M-ficolin (h2): 293T Lysate represents a lysate of human M-ficolin transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.


APPLICATIONS

M-ficolin (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive M-ficolin antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

M-ficolin (036-05-1): sc-80486 is recommended as a positive control antibody for Western Blot analysis of enhanced human M-ficolin expression in M-ficolin transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



M-ficolin (036-05-1): sc-80486. Western blot analysis of M-ficolin expression in non-transfected: sc-117752 (**A**) and human M-ficolin transfected: sc-159506 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.