



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Cytokeratin 18 (h): 293T Lysate: sc-174503

## BACKGROUND

Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation, which is directly applicable to the characterization of malignant tumors. For example, Cytokeratins 10 and 13 are highly expressed in a subset of squamous cell carcinomas while Cytokeratin 18 is expressed in a majority of adenocarcinomas and basal cell carcinomas. Cytokeratin 18 contains two major phosphorylation sites, Ser 33 and Ser 52. Phosphorylation of Ser 18 is essential for the association of Cytokeratin 18 with 14-3-3 proteins and is involved in keratin organization and distribution.

## REFERENCES

- Lauerová, L., Kovarik, J., Bártek, J., Rejthar, A. and Vojtesek, B. 1988. Novel monoclonal antibodies defining epitope of human Cytokeratin 18 molecule. *Hybridoma* 7: 495-504.
- Vojtesek, B., Stasková, Z., Nenutil, R., Lauerová, L., Kovarik, J., Rejthar, A., Bártková, J. and Bártek, J. 1989. Monoclonal antibodies recognizing different epitopes of Cytokeratin 18. *Folia Biol.* 35: 373-382.
- van der Velden, L.A., Schaafsma, H.E., Manni, J.J., Ramaekers, F.C. and Kuijpers, W. 1993. Cytokeratin expression in normal and (pre)malignant head and neck epithelia: an overview. *Head Neck* 15: 133-146.
- Silen, A., Wiklund, B., Norlen, B.J. and Nilsson, S. 1994. Evaluation of a new tumor marker for Cytokeratin 8 and 18 fragments in healthy individuals and prostate cancer patients. *Prostate* 24: 326-332.
- Quillien, V., Ramee, M.P., Bansard, J.Y., Meritte, H., Briens, E., Logeais, Y., Langanay, T., Corbineau, H. and Dazord, L. 1995. Serum and tissue distribution of a fragment of Cytokeratin 19 (CYFRA 21-1) in lung cancer patients. *Anticancer Res.* 15: 2857-2863.
- Marceau, N. and Loranger, A. 1995. Cytokeratin expression, fibrillar organization and subtle function in liver cells. *Biochem. Cell Biol.* 73: 619-625.
- Silen, A., Wiklund, B., Andersson, E.L. and Nilsson, S. 1995. A novel IRMA and ELISA for quantifying Cytokeratin 8 and 18 fragments in the sera of healthy individuals and cancer patients. *Scan. J. Clin. Lab. Invest.* 55: 153-161.
- Mukhopadhyay, T. and Roth, J.A. 1996. Functional inactivation of p53 by antisense RNA induces invasive ability of lung carcinoma cells and down-regulates cytokeatin synthesis. *Anticancer Res.* 16: 1683-1689.
- Ku, N.O., Liao, J. and Omary, M.B. 1998. Phosphorylation of human keratin 18 Serine 33 regulates binding to 14-3-3 proteins. *EMBO J.* 17: 1892-1906.

## CHROMOSOMAL LOCATION

Genetic locus: KRT18 (human) mapping to 12q13.13.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PRODUCT

Cytokeratin 18 (h): 293T Lysate represents a lysate of human Cytokeratin 18 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

Cytokeratin 18 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Cytokeratin 18 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Cytokeratin 18 (DC-10): sc-6259 is recommended as a positive control antibody for Western Blot analysis of enhanced human Cytokeratin 18 expression in Cytokeratin 18 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Cytokeratin 18 (DC-10): sc-6259. Western blot analysis of Cytokeratin 18 expression in non-transfected: sc-117752 (A) and human Cytokeratin 18 transfected: sc-174503 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.