



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# CD68 (h): 293T Lysate: sc-175248

## BACKGROUND

CD68, which is homologous to the mouse antigen macrosialin, belongs to a family of acidic, highly glycosylated lysosomal glycoproteins (LGP) that includes LAMP-1 and LAMP-2. CD68 is found in cytoplasmic granules and in the cytoplasm of various non-hematopoietic tissues including liver and kidney tubules and glomeruli. CD68 is also found, to a lesser extent, on the surface of macrophages, monocytes, neutrophils, basophils and large lymphocytes. LGPs are major components of lysosomal membranes and may act to protect the membranes from attack by hydrolases.

## REFERENCES

1. Fulford, K.A., Sipos, A., Cordell, J.L., Stross, W.P. and Mason, D.Y. 1990. Distribution of the CD68 macrophage/myeloid associated antigen. *Int. Immunol.* 2: 973-980.
2. Fukuda, M. 1991. Lysosomal membrane glycoproteins. Structure, biosynthesis, and intracellular trafficking. *J. Biol. Chem.* 266: 21327-21330.
3. Holness, C.L. and Simmons, D.L. 1993. Molecular cloning of CD68, a human macrophage marker related to lysosomal glycoproteins. *Blood* 81: 1607-1613.
4. Ramprasad, M.P., Fischer, W., Witztum, J.L., Sambrano, G.R., Quehenberger, O. and Steinberg, D. 1995. The 94 to 97 kDa mouse macrophage membrane protein that recognizes oxidized low density lipoprotein and phosphatidylserine-rich liposomes is identical to macrosialin, the mouse homologue of human CD68. *Proc. Natl. Acad. Sci. USA* 92: 9580-9584.
5. Strobl, H., Scheinecker, C., Csmarits, B., Majdic, O. and Knapp, W. 1995. Flow cytometric analysis of intracellular CD68 molecule expression in normal and malignant haemopoiesis. *Br. J. Haematol.* 90: 774-782.
6. Ramprasad, M.P., Terpstra, V., Kondratenko, N., Quehenberger, O. and Steinberg, D. 1996. Cell surface expression of mouse macrosialin and human CD68 and their role as macrophage receptors for oxidized low density lipoprotein. *Proc. Natl. Acad. Sci. USA* 93: 14833-14838.

## CHROMOSOMAL LOCATION

Genetic locus: CD68 (human) mapping to 17p13.1.

## PRODUCT

CD68 (h): 293T Lysate represents a lysate of human CD68 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

CD68 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive CD68 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


CD68 (E-11): sc-17832 is recommended as a positive control antibody for Western Blot analysis of enhanced human CD68 expression in CD68 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG<sub>κ</sub> BP-HRP: sc-516102 or m-IgG<sub>κ</sub> BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



CD68 (E-11): sc-17832. Western blot analysis of CD68 expression in non-transfected: sc-117752 (**A**) and human CD68 transfected: sc-175248 (**B**) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.