



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# IL-8RA (h3): 293T Lysate: sc-176056

## BACKGROUND

IL-8 has been shown to function as a potent neutrophil chemostatic and activating peptide and is an important mediator of inflammatory diseases. Two distinct human IL-8 receptors, designated IL-8RA and IL-8RB, have been characterized. Both are expressed at a high level on neutrophils, and to a lesser extent on monocytes and myeloid cell lines. In addition, the IL-8RA subunit is expressed in T cells such as the Jurkat cell line. Both IL-8Rs are members of the seven-transmembrane domain rhodopsin superfamily of receptors and as such, couple G proteins for signal transduction. The two receptors share 77% amino acid identity. IL-8RA exhibits high affinity binding for IL-8 and low affinity MGSA binding, whereas IL-8RB has high affinity binding for both IL-8 and MGSA.

## REFERENCES

- Holmes, W.E., et al. 1991. Structure and functional expression of a human interleukin-8 receptor. *Science* 253: 1278-1280.
- Murphy, P.M., et al. 1991. Cloning of complementary DNA encoding a functional human interleukin-8 receptor. *Science* 253: 1280-1283.
- Koch, A.E., et al. 1992. Interleukin-8 as a macrophage-derived mediator of angiogenesis. *Science* 258: 1789-1801.
- Lee, J., et al. 1992. Characterization of two high affinity human interleukin-8 receptors. *J. Biol. Chem.* 267: 16283-16287.
- Hebert, C.A., et al. 1993. Interleukin-8: a review. *Cancer Invest.* 11: 743-750.
- Kupper, R.W., et al. 1993. G protein activation by IL-8 and related cytokines in human neutrophil plasma membranes. *Biochem. J.* 282: 429-434.
- Moser, B., et al. 1993. Expression of transcripts for two IL-8 receptors in human phagocytotes, lymphocytes and melanoma cells. *Biochem. J.* 294: 285-292.
- Barnett, M.L., et al. 1993. Characterization of interleukin-8 receptors in human neutrophil membranes: regulation by guanine nucleotides. *Biochim. Biophys. Acta* 1177: 275-282.
- Chunthrapai, A., et al. 1994. Monoclonal antibodies detect different distribution patterns of IL-8 receptor A and IL-8 receptor B on human peripheral blood leukocytes. *J. Immunol.* 153: 5682-5688.

## CHROMOSOMAL LOCATION

Genetic locus: CXCR1 (human) mapping to 2q35.

## PRODUCT

IL-8RA (h3): 293T Lysate represents a lysate of human IL-8RA transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

IL-8RA (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive IL-8RA antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


IL-8RA (B-1): sc-7303. is recommended as a positive control antibody for Western Blot analysis of enhanced human IL-8RA expression in IL-8RA transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG<sub>κ</sub> BP-HRP: sc-516102 or m-IgG<sub>κ</sub> BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



IL-8RA (B-1): sc-7303. Western blot analysis of IL-8RA expression in non-transfected: sc-117752 (**A**) and human IL-8RA transfected: sc-176056 (**B**) 293T whole cell lysates.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.