



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



α -actinin-4 (h): 293T Lysate: sc-176191



BACKGROUND

The spectrin gene family encodes a diverse group of cytoskeletal proteins that include spectrins, dystrophins and α -actinins. There are four tissue-specific α -actinins, namely α -actinin-1, α -actinin-2, α -actinin-3 and α -actinin-4, which are localized to muscle and non-muscle cells, including skeletal, cardiac and smooth muscle cells, as well as within the cytoskeleton. Each α -actinin protein contains one Actin-binding domain, two calponin-homology domains, two EF-hand domains and four spectrin repeats, through which they function as bundling proteins that can cross-link F-Actin, thus anchoring Actin to a variety of intracellular structures. Defects in the gene encoding α -actinin-4 are the cause of focal segmental glomerulosclerosis 1 (FSGS1), a common renal lesion characterized by decreasing kidney function and, ultimately, renal failure.

REFERENCES

1. Youssoufian, H., McAfee, M. and Kwiatkowski, D.J. 1990. Cloning and chromosomal localization of the human cytoskeletal α -actinin gene reveals linkage to the β -spectrin gene. Am. J. Hum. Genet. 47: 62-71.
2. Nishiyama, M., Ozturk, M., Frohlich, M., Mafune, K., Steele, G. and Wands, J.R. 1990. Expression of human α -actinin in human hepatocellular carcinoma. Cancer Res. 50: 6291-6294.
3. Yürük, B. and Niggli, V. 1992. α -actinin and vinculin in human neutrophils: reorganization during adhesion and relation to the Actin network. J. Cell Sci. 101: 403-414.
4. Knudsen, K.A., Soler, A.P., Johnson, K.R. and Wheelock, M.J. 1995. Interaction of α -actinin with the cadherin/catenin cell-cell adhesion complex via α -catenin. J. Cell Biol. 130: 67-77.
5. Reinhard, M., Zumbrunn, J., Jaquemar, D., Kuhn, M., Walter, U. and Trueb, B. 1999. An α -actinin binding site of Zyxin is essential for subcellular Zyxin localization and α -actinin recruitment. J. Biol. Chem. 274: 13410-13418.
6. Harper, B.D., Beckerle, M.C. and Pomiès, P. 2000. Fine mapping of the α -actinin binding site within cysteine-rich protein. Biochem. J. 350: 269-274.
7. Gonzalez, A.M., Otey, C., Edlund, M. and Jones, J.C. 2001. Interactions of a hemidesmosome component and actinin family members. J. Cell Sci. 114: 4197-4206.
8. Bois, P.R., Borgon, R.A., Vonrhein, C. and Izard, T. 2005. Structural dynamics of α -actinin-vinculin interactions. Mol. Cell. Biol. 25: 6112-6122.
9. Nyman-Huttunen, H., Tian, L., Ning, L. and Gahmberg, C.G. 2006. α -actinin-dependent cytoskeletal anchorage is important for ICAM-5-mediated neuritic outgrowth. J. Cell Sci. 119: 3057-3066.

CHROMOSOMAL LOCATION

Genetic locus: ACTN4 (human) mapping to 19q13.2.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PRODUCT

α -actinin-4 (h): 293T Lysate represents a lysate of human α -actinin-4 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

α -actinin-4 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive α -actinin-4 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


α -actinin-4 (B-11): sc-390180 is recommended as a positive control antibody for Western Blot analysis of enhanced human α -actinin-4 expression in α -actinin-4 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



α -actinin-4 (B-11): sc-390180. Western blot analysis of α -actinin-4 expression in non-transfected: sc-117752 (A) and human α -actinin-4 transfected: sc-176191 (B) 293T whole cell lysates.

α -actinin-4 (B-11): sc-390180. Western blot analysis of α -actinin-4 expression in non-transfected: sc-117752 (A) and human α -actinin-4 transfected: sc-176191 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.