



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PTP IA-2 β (h): 293T Lysate: sc-176447

BACKGROUND

Protein-Tyrosine Phosphatase Receptor-Type IA-2 β (PTP IA-2 β), alternately known as PTPRN2 or Phogrin, localizes in dense-core secretory vesicles of pancreas islet cells and influences Insulin secretion. The PTP IA-2 β precursor is an autoantigen that contributes to Insulin-dependent diabetes mellitus (IDDM). The autoantigenic epitopes of PTP IA-2 β appear within the cytoplasmic domain of this transmembrane protein. PTP IA-2 β is present at high levels in brain and pancreas with lower levels in trachea, prostate, stomach and spinal cord. The human PTPRN2 gene maps to chromosome 7q36.3. Northern blot analysis showed that PTPRN2 was expressed as 5.5- and 3.7-kb transcripts primarily in human brain and pancreas. Three alternative transcript splice variants of this gene encode distinct proteins.

REFERENCES

1. Kawasaki, E., et al. 1996. Molecular cloning and characterization of the human transmembrane protein tyrosine phosphatase homologue, phogrin, an autoantigen of type 1 diabetes. *Biochem. Biophys. Res. Commun.* 227: 440-447.
2. Smith, P.D., et al. 1996. ICAAR, a novel member of a new family of transmembrane, tyrosine phosphatase-like proteins. *Biochem. Biophys. Res. Commun.* 229: 402-411.
3. Kubosaki, A., et al. 2004. Targeted disruption of the IA-2 β gene causes glucose intolerance and impairs Insulin secretion but does not prevent the development of diabetes in NOD mice. *Diabetes* 53: 1684-1691.
4. Drake, P.G., et al. 2003. A novel strategy for the development of selective active-site inhibitors of the protein tyrosine phosphatase-like proteins islet-cell antigen 512 (IA-2) and phogrin (IA-2 β). *Biochem. J.* 373: 393-401.
5. Achenbach, P., et al. 2002. Spontaneous peripheral T cell responses to the IA-2 β (phogrin) autoantigen in young nonobese diabetic mice. *J. Autoimmun.* 19: 111-116.
6. Gross, S., et al. 2002. Multimerization of the protein-tyrosine phosphatase (PTP)-like Insulin-dependent diabetes mellitus autoantigens IA-2 and IA-2 β with receptor PTPs (RPTPs). Inhibition of RPTP α enzymatic activity. *J. Biol. Chem.* 277: 48139-48145.
7. SWISS-PROT/TrEMBL (Q92932). World Wide Web URL:
<http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: PTPRN2 (human) mapping to 7q36.3.

PRODUCT

PTP IA-2 β (h): 293T Lysate represents a lysate of human PTP IA-2 β transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

PTP IA-2 β (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive PTP IA-2 β antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.