



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# Mcl-1 (h): 293T Lysate: sc-176626

## BACKGROUND

B cell CLL/lymphoma-2 (Bcl-2) blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal. Myeloid cell leukemia-1 (Mcl-1) shares sequence homology with Bcl-2 and further resembles Bcl-2 in that its expression promotes cell viability. p53 and Mcl-1 demonstrate opposing effects on mitochondrial apoptosis by mediating Bcl-2 antagonist killer (Bak) activity. Mcl-1 is an important and specific regulator that is necessary for the homeostasis of early hematopoietic progenitors. Glycogen synthase kinase-3 (GSK-3) controls Mcl-1 stability, which has an effect on the regulation of apoptosis by growth factors PI 3-kinase and Akt. Mice with a deficiency of the Mcl-1 protein show a significant reduction in B and T lymphocytes similar to the effects observed in IL-7- or IL-7R-deficient mice.

## REFERENCES

- Kozopas, K.M., et al. 1993. Mcl-1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to Bcl-2. Proc. Natl. Acad. Sci. USA 90: 3516-3520.
- Craig, R.W., et al. 1994. Human and mouse chromosomal mapping of the myeloid cell leukemia-1 gene: Mcl-1 maps to human chromosome 1q21, a region that is frequently altered in preneoplastic and neoplastic disease. Genomics 23: 457-463.
- Rinkenberger, J.L., et al. 2000. Mcl-1 deficiency results in peri-implantation embryonic lethality. Genes Dev. 14: 23-27.
- Bae, J., et al. 2000. Mcl-1S, a splicing variant of the anti-apoptotic Bcl-2 family member Mcl-1, encodes a pro-apoptotic protein possessing only the BH3 domain. J. Biol. Chem. 275: 25255-25261.
- Opferman, J.T., et al. 2003. Development and maintenance of B and T lymphocytes requires anti-apoptotic Mcl-1. Nature 426: 671-676.
- Leu, J.I., et al. 2004. Mitochondrial p53 activates Bak and causes disruption of a Bak-Mcl-1 complex. Nat. Cell Biol. 6: 443-450.
- Opferman, J.T., et al. 2005. Obligate role of anti-apoptotic Mcl-1 in the survival of hematopoietic stem cells. Science 307: 1101-1104.
- Maurer, U., et al. 2006. Glycogen synthase kinase-3 regulates mitochondrial outer membrane permeabilization and apoptosis by destabilization of Mcl-1. Mol. Cell 21: 749-760.

## CHROMOSOMAL LOCATION

Genetic locus: MCL1 (human) mapping to 1q21.3.

## PRODUCT

Mcl-1 (h): 293T Lysate represents a lysate of human Mcl-1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

Mcl-1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Mcl-1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Mcl-1 (C-2): sc-377487 is recommended as a positive control antibody for Western Blot analysis of enhanced human Mcl-1 expression in Mcl-1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgG<sub>x</sub> BP-HRP: sc-516102 or m-IgG<sub>x</sub> BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Mcl-1 (C-2): sc-377487. Western blot analysis of Mcl-1 expression in non-transfected: sc-117752 (**A**) and human Mcl-1 transfected: sc-176626 (**B**) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.