



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

V-ATPase D1 (h): 293T Lysate: sc-176665

BACKGROUND

Vacuolar-type H⁺-ATPase (V-ATPase) is a multisubunit enzyme responsible for the acidification of eukaryotic intracellular organelles. V-ATPases pump protons against an electrochemical gradient, while F-ATPases reverse the process, thereby synthesizing ATP. A peripheral V₁ domain, which is responsible for ATP hydrolysis, and an integral V₀ domain, which is responsible for proton translocation, comprise the V-ATPase complex. Nine subunits (A-H) make up the V₁ domain and five subunits (A, D, C, C' and C'') make up the V₀ domain. V-ATPase D1 (ATPase, H⁺ transporting, lysosomal, V₀ subunit D1), also known as ATP6V0D1, P39, VATX, VMA6, ATP6D or VPATPD, is the D subunit of the V₀ domain. Expressed ubiquitously, V-ATPase D1 acts in concert with other V₀ subunits to catalytically acidify a variety of intracellular compartments, thereby synthesizing ATP to be used for vacuolar transport.

REFERENCES

- van Hille, B., Vanek, M., Richener, H., Green, J.R. and Bilbe, G. 1993. Cloning and tissue distribution of subunits C, D, and E of the human vacuolar H⁺-ATPase. *Biochem. Biophys. Res. Commun.* 197: 15-21.
- Finbow, M.E. and Harrison, M.A. 1997. The vacuolar H⁺-ATPase: a universal proton pump of eukaryotes. *Biochem. J.* 324: 697-712.
- Forgac, M. 1999. Structure and properties of the vacuolar (H⁺)-ATPases. *J. Biol. Chem.* 274: 12951-12954.
- Agarwal, A.K. and White, P.C. 2000. Structure of the VPATPD gene encoding subunit D of the human vacuolar proton ATPase. *Biochem. Biophys. Res. Commun.* 279: 543-547.
- Smith, A.N., Borthwick, K.J. and Karet, F.E. 2002. Molecular cloning and characterization of novel tissue-specific isoforms of the human vacuolar H⁺-ATPase C, G and d subunits, and their evaluation in autosomal recessive distal renal tubular acidosis. *Gene* 297: 169-177.
- Nishi, T. and Forgac, M. 2002. The vacuolar (H⁺)-ATPases—nature's most versatile proton pumps. *Nat. Rev. Mol. Cell Biol.* 3: 94-103.
- Morel, N. 2003. Neurotransmitter release: the dark side of the vacuolar-H⁺-ATPases. *Biol. Cell.* 95: 453-457.
- Kawasaki-Nishi, S., Nishi, T. and Forgac, M. 2003. Proton translocation driven by ATP hydrolysis in V-ATPases. *FEBS Lett.* 545: 76-85.
- Pietrement, C., Sun-Wada, G.H., Silva, N.D., McKee, M., Marshansky, V., Brown, D., Futai, M. and Breton, S. 2006. Distinct expression patterns of different subunit isoforms of the V-ATPase in the rat epididymis. *Biol. Reprod.* 74: 185-194.

CHROMOSOMAL LOCATION

Genetic locus: ATP6V0D1 (human) mapping to 16q22.1.

PRODUCT

V-ATPase D1 (h): 293T Lysate represents a lysate of human V-ATPase D1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

V-ATPase D1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive V-ATPase D1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

V-ATPase D1 (34-Z): sc-81887 is recommended as a positive control antibody for Western Blot analysis of enhanced human V-ATPase D1 expression in V-ATPase D1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



V-ATPase D1 (34-Z): sc-81887. Western blot analysis of V-ATPase D1 expression in non-transfected: sc-117752 (A) and human V-ATPase D1 transfected: sc-176665 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.