



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



IL-1 β (h): 293T Lysate: sc-176712

BACKGROUND

Two forms of interleukin-1, designated IL-1 α and IL-1 β , have been described. Although encoded by distinct genes and exhibiting roughly only 25% sequence identity, IL-1 α and IL-1 β bind to the same receptor and seem to elicit similar biological responses. IL-1 production is generally thought to be associated with inflammation, but it has also been shown to be expressed during kidney development, thymocyte differentiation and cartilage degradation. IL-1 plays a critical role in the regulation of immune response and inflammation, acting as an activator of T and B lymphocytes and natural killer (NK) cells. In T cells, IL-1 stimulates the production of IL-2 and selectively inhibits IL-4 expression. IL-1 induces B cell proliferation and maturation, and immunoglobulin synthesis. NK cells require IL-1 β for production of the anti-pathogen IFN- γ . IL-1 has also been implicated in several pathological conditions including rheumatoid arthritis, inflammatory bowel disease and atherosclerosis.

REFERENCES

- Auron, P.E., Webb, A.C., Rosenwasser, L.J., Mucci, S.F., Rich, A., Wolff, S.M. and Dinarello, C.A. 1985. Nucleotide sequence of human monocyte interleukin-1 precursor cDNA. Proc. Natl. Acad. Sci. USA 81: 7907-7911.
- March, C.J., Mosley, B., Larsen, A., Cerretti, D.P., Braedt, G., Price, V., Gillis, S., Henney, C.S., Kronheim, S.R. and Grabstein, K. 1985. Cloning, sequence and expression of two distinct human interleukin-1 complementary DNAs. Nature 315: 641-647.
- Dinarello, C.A. 1991. Interleukin-1 and interleukin-1 antagonism. Blood 77: 1627-1652.
- Sadouk, M.B., Pelletier, J.P., Tardif, G., Kiansa, K., Cloutier, J.M. and Martel-Pelletier, J. 1995. Human synovial fibroblasts coexpress IL-1 receptor type I and type II mRNA. The increased level of the IL-1 receptor in osteoarthritis cells is related to an increased level of the type I receptor. Lab. Invest. 73: 347-355.
- Lonnemann, G., Shapiro, L., Engler-Blum, G., Müller, G.A., Koch, K.M. and Dinarello, C.A. 1995. Cytokines in human renal interstitial fibrosis. I. Interleukin-1 is a paracrine growth factor for cultured fibrosis-derived kidney fibroblasts. Kidney Int. 47: 837-844.
- Zúñiga-Pflücker, J.C., Di, J. and Lenardo, M.J. 1995. Requirement for TNF α and IL-1 α in fetal thymocyte commitment and differentiation. Science 268: 1906-1909.
- Sandborg, C.I., Imfeld, K.L., Zaldivar, F., Wang, Z., Buckingham, B.A. and Berman, M.A. 1995. IL-4 expression in human T cells is selectively inhibited by IL-1 α and IL-1 β . J. Immunol. 155: 5206-5212.

CHROMOSOMAL LOCATION

Genetic locus: IL1B (human) mapping to 2q13.

PRODUCT

IL-1 β (h): 293T Lysate represents a lysate of human IL-1 β transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

IL-1 β (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive IL-1 β antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


IL-1 β (E-12): sc-515787 is recommended as a positive control antibody for Western Blot analysis of enhanced human IL-1 β expression in IL-1 β transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



IL-1 β (E-12): sc-515787. Western blot analysis of IL-1 β expression in non-transfected: sc-117752 (**A**) and human IL-1 β transfected: sc-176712 (**B**) 293T whole cell lysates.

IL-1 β (E7-2-hIL β): sc-32294. Western blot analysis of IL-1 β expression in non-transfected: sc-117752 (**A**) and human IL-1 β transfected: sc-176712 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.