



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



GADD 153 (h): 293T Lysate: sc-177265

BACKGROUND

GADD 153 has been described as a growth arrest and DNA damage-inducible gene that encodes a C/EBP-related nuclear protein. This protein has also been designated C/EBP-homologous protein (CHOP-10). GADD 153 expression is induced by a variety of cellular stresses, inducing nutrient deprivation and metabolic perturbations. GADD 153 functions to block cells in G₁ to S phase in cell cycle progression and acts by dimerizing with other C/EBP proteins to direct GADD 153 dimers away from "classical" C/EBP binding sites, recognizing instead unique "nonclassical" sites. Thus GADD 153 acts as a negative modulator of C/EBP-like proteins in certain terminally differentiated cells, similar to the regulatory function of Id on the activity of Myo D and Myo D-related proteins involved in the development of muscle cells.

REFERENCES

- Sherr, C.J. 1994. G₁ phase progression: cycling on cue. *Cell* 79: 551-555.
- Hunter, T., et al. 1994. Cyclins and cancer II: cyclin D and CDK inhibitors come of age. *Cell* 79: 573-582.
- Ron, D. 1994. Inducible growth arrest: new mechanistic insights. *Proc. Natl. Acad. Sci. USA* 91: 1985-1986.
- Smith, M.L., et al. 1994. Interaction of the p53-regulated protein GADD 45 with proliferating cell nuclear antigen. *Science* 266: 1376-1380.
- Gujuluva, C.N., et al. 1994. Effect of UV-irradiation on cell cycle, viability and the expression of p53, GADD 153 and GADD 45 genes in normal and HPV-immortalized human oral keratinocytes. *Oncogene* 9: 1819-1827.
- Selvakumaran, M., et al. 1994. The novel primary response gene MyD118 and the proto-oncogenes Myb, Myc, and Bcl-2 modulate transforming growth factor β 1-induced apoptosis of myeloid leukemia cells. *Mol. Cell. Biol.* 14: 2352-2360.
- Zhan, Q., et al. 1994. The GADD and MyD genes define a novel set of mammalian genes encoding acidic proteins that synergistically suppress cell growth. *Mol. Cell. Biol.* 14: 2361-2371.
- Su, Z.Z., et al. 1997. Subtraction hybridization identifies a transformation progression associated-gene PEG-3 with sequence homology to a growth arrest and DNA damage-inducible gene. *Proc. Natl. Acad. Sci. USA* 94: 9125-9130.
- Ito, A., et al. 2000. Bystander-killing effect and cyclic induction of TNF α gene under heat-inducible promoter GADD 153. *J. Biosci. Bioeng.* 90: 437-441.

CHROMOSOMAL LOCATION

Genetic locus: DDIT3 (human) mapping to 12q13.3.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

GADD 153 (h): 293T Lysate represents a lysate of human GADD 153 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

GADD 153 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive GADD 153 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

GADD 153 (B-3): sc-7351 is recommended as a positive control antibody for Western Blot analysis of enhanced human GADD 153 expression in GADD 153 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



GADD 153 (B-3): sc-7351. Western blot analysis of GADD 153 expression in non-transfected: sc-117752 (**A**) and human GADD 153 transfected: sc-177265 (**B**) 293T whole cell lysates.

GADD 153 (6D313): sc-71136. Western blot analysis of GADD 153 expression in non-transfected: sc-117752 (**A**) and human GADD 153 transfected: sc-177265 (**B**) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.