



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Mel-CAM (h2): 293T Lysate: sc-177535

BACKGROUND

The tumorigenic and metastatic phenotype of melanoma cells correlates well with an increased expression of cell-cell and cell-matrix adhesion receptors. The human Mel-CAM gene encodes a transmembrane glycoprotein, also designated MCAM, MUC18 or CD146, that belongs to the immunoglobulin superfamily and functions as a Ca^{2+} -independent cell adhesion molecule. The deduced human sequence of 603 amino acids consists of a signal peptide, 5 immunoglobulin-like domains, a transmembrane region and a short cytoplasmic tail. Mel-CAM expression is restricted to advanced primary and metastatic melanomas and to cell lines of the neuroectodermal lineage, but not normal melanocytes. Mel-CAM is found on 80% of advanced primary human melanomas and correlates well with development of metastatic disease. Mel-CAM activation initiates an outside-in signaling pathway that involves the protein tyrosine kinases Fyn, FAK and paxillin. Mel-CAM influences the dynamics of actin cytoskeleton rearrangement and is essential for the maintenance of thymic architecture and function.

REFERENCES

- Lehmann, J.M., et al. 1989. MUC18, a marker of tumor progression in human melanoma, shows sequence similarity to the neural cell adhesion molecules of the immunoglobulin superfamily. *Proc. Natl. Acad. Sci. USA* 86: 9891-9895.
- Kuzu, I., et al. 1993. Expression of adhesion molecules on the endothelium of normal tissue vessels and vascular tumors. *Lab. Invest.* 69: 322-328.
- Sers, C., et al. 1993. Genomic organization of the melanoma-associated glycoprotein MUC18: implications for the evolution of the immunoglobulin domains. *Proc. Natl. Acad. Sci. USA* 90: 8514-8518.
- Shih, I.M. 1999. The role of CD146 (Mel-CAM) in biology and pathology. *J. Pathol.* 189: 4-11.
- Seftalioglu, A. and Karakoc, L. 2000. Expression of CD146 adhesion molecules (MUC18 or MCAM) in the thymic microenvironment. *Acta Histochem.* 102: 69-83.
- Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 155735. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Satyamoorthy, K., et al. 2001. Mel-CAM-specific genetic suppressor elements inhibit melanoma growth and invasion through loss of gap junctional communication. *Oncogene* 20: 4676-4684.

CHROMOSOMAL LOCATION

Genetic locus: MCAM (human) mapping to 11q23.3.

PRODUCT

Mel-CAM (h2): 293T Lysate represents a lysate of human Mel-CAM transfected 293T cells and is provided as 100 μg protein in 200 μl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Mel-CAM (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Mel-CAM antibodies. Recommended use: 10-20 μl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


Mel-CAM (A-9): sc-374556 is recommended as a positive control antibody for Western Blot analysis of enhanced human Mel-CAM expression in Mel-CAM transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Mel-CAM (A-9): sc-374556. Western blot analysis of Mel-CAM expression in non-transfected: sc-117752 (**A**) and human Mel-CAM transfected: sc-177535 (**B**) 293T whole cell lysates.

Mel-CAM (B-10): sc-376762. Western blot analysis of Mel-CAM expression in non-transfected: sc-117752 (**A**) and human Mel-CAM transfected: sc-177535 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.