



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ZAP-70 (G-4): sc-17760

BACKGROUND

The activation of T lymphocytes by antigens is mediated by the T cell receptor (TCR) which is a multisubunit complex assembled from at least six different genes. The TCR subunits include the T α and β chains, the CD3 γ , δ and ϵ chains and a ζ -containing homodimer or heterodimer. The disulfide-linked T α - β heterodimer is responsible for antigen recognition, but the short 5 amino acid cytoplasmic domains of T α and β are unlikely to be sufficient to couple to intracellular signaling pathways. In contrast, the structured features of the CD3 and ζ subunits suggest a role in signal transduction. Of these, the ζ chain, which is expressed as either a homodimer or heterodimer, has a short extracellular domain of only 9 amino acids, but a larger 113 amino acid cytoplasmic domain. A 70 kDa tyrosine phosphoprotein, ZAP-70, has been identified that associates with ζ and undergoes tyrosine phosphorylation following TCR stimulation.

REFERENCES

1. Clevers, H., et al. 1988. The T cell receptor/CD3 complex: a dynamic protein ensemble. *Annu. Rev. Immunol.* 6: 629-662.
2. Baniyash, M., et al. 1988. Disulfide linkage of the ζ and η chains of the T cell receptor. *J. Biol. Chem.* 263: 9874-9878.
3. Baniyash, M., et al. 1988. The T cell antigen receptor ζ chain is tyrosine phosphorylated upon activation. *J. Biol. Chem.* 263: 18225-18230.
4. Baniyash, M., et al. 1989. The isolation and characterization of the murine T cell antigen receptor ζ chain gene. *J. Biol. Chem.* 264: 13252-13257.
5. Frank, S.J., et al. 1990. The structure and signaling function of the invariant T cell receptor components. *Semin. Immunol.* 2: 89-97.
6. Clayton, L.K., et al. 1991. CD3 η and CD3 ζ are alternatively spliced products of a common genetic locus and are transcriptionally and/or post-transcriptionally regulated during T-cell development. *Proc. Natl. Acad. Sci. USA* 88: 5202-5206.

CHROMOSOMAL LOCATION

Genetic locus: ZAP70 (human) mapping to 2q11.2.

SOURCE

ZAP-70 (G-4) is a mouse monoclonal antibody raised against amino acids 253-304 of ZAP-70 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

ZAP-70 (G-4) is available conjugated to phycoerythrin (sc-17760 PE), 200 μ g/ml, for IF, IHC(P) and FCM.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS


ZAP-70 (G-4) is recommended for detection of ZAP-70 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1,000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), flow cytometry (1 μ g per 1 x 10⁶ cells) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ZAP-70 siRNA (h): sc-29526, ZAP-70 shRNA Plasmid (h): sc-29526-SH and ZAP-70 shRNA (h) Lentiviral Particles: sc-29526-V.


Molecular Weight of ZAP-70: 70 kDa.

Positive Controls: CCRF-CEM cell lysate: sc-2225, Jurkat whole cell lysate: sc-2204 or MOLT-4 cell lysate: sc-2233.

DATA



ZAP-70 (G-4): sc-17760. Western blot analysis of ZAP-70 expression in Jurkat (A) and MOLT-4 (B) whole cell lysates.



ZAP-70 (G-4) PE: sc-17760 PE. Intracellular FCM analysis of fixed and permeabilized Jurkat cells. Black line histogram represents the isotype control, normal mouse IgG₁: sc-2866.

SELECT PRODUCT CITATIONS

1. Kondo, T., et al. 2001. Dendritic cells signal T cells in the absence of exogenous antigen. *Nat. Immunol.* 2: 932-938.
2. Zhang, W., et al. 2003. Negative regulation of T cell antigen receptor-mediated Crk-L-C3G signaling and cell adhesion by Cbl- β . *J. Biol. Chem.* 278: 23978-23983.
3. Slack, G.W., et al. 2007. Flow cytometric detection of ZAP-70 in chronic lymphocytic leukemia: correlation with immunocytochemistry and Western blot analysis. *Arch. Pathol. Lab. Med.* 131: 50-56.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **ZAP-70 (1E7.2): sc-32760** for ZAP-70 antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647.