



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



ZNF276 (h4): 293T Lysate: sc-178164

BACKGROUND

Zinc finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZNF276 is a 614 amino acid protein containing 5 C₂H₂-type zinc fingers and one zinc finger associated (ZAD) domain. Due to a loss of heterozygosity at the chromosomal location of the gene encoding ZNF276 in sporadic breast cancers, the ZNF276 gene has been targeted as a possible breast cancer tumor suppressor. The FANCA gene, which encodes a DNA repair protein, is situated at the same chromosomal location as the ZNF276 gene, suggesting a possible involvement of ZNF276 in the progression of Fanconi anemia, an autosomal recessive disorder which is caused by mutations in the gene encoding FANCA. There are two isoforms of ZNF276 that exist as a result of an alternative splicing event.

REFERENCES

1. Payre, F. and Vincent, A. 1988. Finger proteins and DNA-specific recognition: distinct patterns of conserved amino acids suggest different evolutionary modes. FEBS Lett. 234: 245-250.
2. Brenner, A.J. and Aldaz, C.M. 1997. The genetics of sporadic breast cancer. Prog. Clin. Biol. Res. 396: 63-82.
3. Nakamura, A., Matsuura, S., Tauchi, H., Hanada, R., Ohashi, H., Hasegawa, T., Honda, K., Masuno, M., Imaizumi, K., Sugita, K., Ide, T. and Komatsu, K. 1999. Four novel mutations of the Fanconi anemia group A gene (FAA) in Japanese patients. J. Hum. Genet. 44: 48-51.
4. Wong, J.C., Alon, N., Norga, K., Kruty, F.A., Youssoufian, H. and Buchwald, M. 2000. Cloning and analysis of the mouse Fanconi anemia group A cDNA and an overlapping penta zinc finger cDNA. Genomics 67: 273-283.
5. Chung, H.R., Schäfer, U., Jäckle, H. and Böhm, S. 2002. Genomic expansion and clustering of ZAD-containing C₂H₂ zinc finger genes in *Drosophila*. EMBO Rep. 3: 1158-1162.
6. Wong, J.C., Gokgoz, N., Alon, N., Andrusis, I.L. and Buchwald, M. 2003. Cloning and mutation analysis of ZFP276 as a candidate tumor suppressor in breast cancer. J. Hum. Genet. 48: 668-671.
7. Jauch, R., Bourenkov, G.P., Chung, H.R., Urlaub, H., Reidt, U., Jäckle, H. and Wahl, M.C. 2003. The zinc finger-associated domain of the *Drosophila* transcription factor grauzone is a novel zinc-coordinating protein-protein interaction module. Structure 11: 1393-1402.
8. Imami, K., Sugiyama, N., Kyono, Y., Tomita, M. and Ishihama, Y. 2008. Automated phosphoproteome analysis for cultured cancer cells by two-dimensional nanoLC-MS using a calcined titania/C18 biphasic column. Anal. Sci. 24: 161-166.
9. Liu, J. and Stormo, G.D. 2008. Context-dependent DNA recognition code for C₂H₂ zinc finger transcription factors. Bioinformatics 24: 1850-1857.

CHROMOSOMAL LOCATION

Genetic locus: ZNF276 (human) mapping to 16q24.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

ZNF276 (h4): 293T Lysate represents a lysate of human ZNF276 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

ZNF276 (h4): 293T Lysate is suitable as a Western Blotting positive control for human reactive ZNF276 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


ZNF276 (F-10): sc-398207 is recommended as a positive control antibody for Western Blot analysis of enhanced human ZNF276 expression in ZNF276 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG_κ BP-HRP: sc-516102 or m-IgG_κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



ZNF276 (F-10): sc-398207. Western blot analysis of ZNF276 expression in non-transfected: sc-117752 (**A**) and human ZNF276 transfected: sc-178164 (**B**) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.