



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



CD209b (m): 293 Lysate: sc-178374

BACKGROUND

Antigen-presenting cells are localized in essentially every tissue, where they operate at the interface of innate and acquired immunity by capturing pathogens and presenting pathogen-derived peptides to T cells. Dendritic cells capture antigens or viruses in peripheral tissue and transport them to lymphoid organs, an event that induces cellular T cell responses. The mouse CD209 family of cell adhesion receptors consists of CD209a (also known as DC-SIGN), CD209b, CD209c, CD209d, CD209e, CD209f and CD209g, all of which function to mediate the endocytosis and subsequent degradation of pathogens within lysosomal compartments. There are two human CD209 proteins, designated DC-SIGN and DC-SIGNR, which function in a similar manner to the mouse proteins.

REFERENCES

1. Engering A., et al. 2002. The dendritic cell-specific adhesion receptor DC-SIGN internalizes antigen for presentation to T cells. *J. Immunol.* 168: 2118-2126.
2. Geijtenbeek T.B., et al. 2002. Marginal zone macrophages express a murine homologue of DC-SIGN that captures blood-borne antigens *in vivo*. *Blood* 100: 2908-2916.
3. Moris A., et al. 2004. DC-SIGN promotes exogenous MHC I-restricted HIV-1 antigen presentation. *Blood* 103: 2648-2654.
4. Cormier, E.G., et al. 2004. L-SIGN (CD209L) and DC-SIGN (CD209) mediate transinfection of liver cells by hepatitis C virus. *Proc. Natl. Acad. Sci. USA* 101: 14067-14072.
5. Weber, K.S., et al. 2004. Sialylation of ICAM-2 on platelets impairs adhesion of leukocytes via LFA-1 and DC-SIGN. *Inflammation* 28: 177-188.
6. Sakuntabhai, A., et al. 2005. A variant in the CD209 promoter is associated with severity of dengue disease. *Nat. Genet.* 37: 507-513.
7. de la Rosa, G., et al. 2005. Regulated recruitment of DC-SIGN to cell-cell contact regions during zymosan-induced human dendritic cell aggregation. *J. Leukoc. Biol.* 77: 699-709.
8. Groot, F., et al. 2005. Lactoferrin prevents dendritic cell-mediated human immunodeficiency virus type 1 transmission by blocking the DC-SIGN-gp120 interaction. *J. Virol.* 79: 3009-3015.
9. Davis, C.W., et al. 2006. The location of asparagine-linked glycans on West Nile virions controls their interactions with CD209 (dendritic cell-specific ICAM-3 grabbing nonintegrin). *J. Biol. Chem.* 281: 37183-37194.

CHROMOSOMAL LOCATION

Genetic locus: Cd209b (mouse) mapping to 8 A1.1.

PRODUCT

CD209b (m): 293 Lysate represents a lysate of mouse CD209b transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CD209b (m): 293 Lysate is suitable as a Western Blotting positive control for mouse reactive CD209b antibodies. Recommended use: 10-20 µl per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.