



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DPM2 siRNA(m): sc-41512

BACKGROUND

Biosynthesis of glycosylphosphatidylinositol and N-glycan pre-cursor is dependent upon a mannosyl donor, dolichol phosphate-mannose (DPM). DPM synthase, a transmembrane protein, is associated with membranes of the rough endoplasmic reticulum and catalyzes mannosyl transfer from GDP-mannose hydrophobic long-chain acceptor dolichyl-phosphate. DPM synthase in various organisms are grouped into two types. One type is a single-component enzyme, represented by *Saccharomyces cerevisiae*, and the other is a multicomponent enzyme which is represented by human DPM synthase and consists of three subunits: DPM1, DPM2 and DPM3. DPM1 is not sufficient for DPM synthesis, which requires the 84 amino acid DPM2 protein for localization to the ER and stable expression of DPM1. The third subunit, DPM3, comprises 92 amino acids, and it is associated with DPM1 via its C-terminal domain and with DPM2 via its N-terminal region. The stability of DPM1 is directly dependent upon DPM3, which is stabilized by DPM2. DPM synthase activity is associated with an ER phosphoprotein. In addition, a mitochondrial DPM synthase exists, which is located on the cytosolic face of the outer membrane of mitochondria.

REFERENCES

1. Gasnier, F., et al. 1992. Mitochondrial dolichyl-phosphate mannose synthase. Purification and immunogold localization by electron microscopy. *Eur. J. Biochem.* 206: 853-858.
2. Forsee, W.T., et al. 1997. Characterization of recombinant yeast dolichyl mannosyl phosphate synthase and site-directed mutagenesis of its cysteine residues. *Eur. J. Biochem.* 244: 935-938.
3. Maeda, Y., et al. 1998. DPM2 regulates biosynthesis of dolichol phosphate-mannose in mammalian cells: correct subcellular localization and stabilization of DPM1, and binding of dolichol phosphate. *EMBO J.* 17: 4920-4929.
4. Tomita, S., et al. 1998. A homologue of *Saccharomyces cerevisiae* Dpm1p is not sufficient for synthesis of dolichol-phosphate-mannose in mammalian cells. *J. Biol. Chem.* 273: 9249-9254.
5. Banerjee, D.K., et al. 1999. Mannosylphosphodolichol synthase activity is associated with a 32 kDa phosphoprotein. *Biosci. Rep.* 19: 169-177.

CHROMOSOMAL LOCATION

Genetic locus: Dpm2 (mouse) mapping to 2 B.

PRODUCT

DPM2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DPM2 shRNA Plasmid (m): sc-41512-SH and DPM2 shRNA (m) Lentiviral Particles: sc-41512-V as alternate gene silencing products.

For independent verification of DPM2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41512A, sc-41512B and sc-41512C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DPM2 siRNA (m) is recommended for the inhibition of DPM2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DPM2 gene expression knockdown using RT-PCR Primer: DPM2 (m)-PR: sc-41512-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.