

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# CIB shRNA (m) Lentiviral Particles: sc-43272-V



The Power to Question

#### **BACKGROUND**

Platelets regulate the function of Integrin  $\alpha 2b/\beta 3$  (GPIIb/IIIa), the platelet Fibrinogen receptor, which is involved in the binding of proteins to integrin cytoplasmic domains. A novel protein, CIB, for calcium- and integrin-binding protein (also designated as Kip for kinase interacting protein, SIP2-28 and DNA-PK\_{cs} interacting protein), binds specifically at the cytoplasmic domain of  $\alpha 2b$  by a number of positively charged residues in its binding site. Binding of CIB to the  $\alpha 2b$  is affected by fluctuations in the intracellular calcium concentration. In aggregated platelets, endogenous CIB and  $\alpha 2b/\beta 3$  translocate to the Triton X-100-insoluble cytoskeleton, demonstrating that the cellular localization of CIB is regulated. CIB also binds to DNA-PK\_{cs}, which is a nuclear protein serine/threonine kinase that plays a role in the DNA repair and recombination process of lymphoid development. Fnk also binds to the CIB, suggesting that CIB may be a regulatory subunit of polo-like kinases. CIB shows significant homology to calcineurin B and calmodulin, and its mRNA levels are ubiquitously expressed in various human tissues.

#### **REFERENCES**

- 1. Naik, U.P., et al. 1997. Identification of a novel calcium-binding protein that interacts with the Integrin  $\alpha$ 2b cytoplasmic domain. J. Biol. Chem. 272: 4651-4654.
- Wu, X., et al. 1997. Interaction between DNA-dependent protein kinase and a novel protein, Kip. Mutat. Res. 385: 13-20.
- 3. Shock, D.D., et al. 1999. Calcium-dependent properties of CIB binding to the Integrin  $\alpha$ 2b cytoplasmic domain and translocation to the platelet cytoskeleton. Biochem. J. 342: 729-735.
- Seki, N., et al. 1999. Structure, expression profile and chromosomal location of an isolog of DNA-PK<sub>cs</sub> interacting protein (Kip) gene. Biochim. Biophys. Acta 1444: 143-147.
- 5. Hwang, P.M., et al. 2000. Structures of the platelet calcium- and integrinbinding protein and the Integrin  $\alpha$ 2b cytoplasmic domain suggest a mechanism for calcium-regulated recognition; homology modeling and NMR studies. J. Mol. Recognit. 13: 83-92.

#### CHROMOSOMAL LOCATION

Genetic locus: Cib1 (mouse) mapping to 7 D3.

#### **PRODUCT**

CIB shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200  $\mu$ l frozen stock containing 1.0 x 10<sup>6</sup> infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see CIB siRNA (m): sc-43272 and CIB shRNA Plasmid (m): sc-43272-SH as alternate gene silencing products.

#### **STORAGE**

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at  $4^\circ$  C for up to one week. Avoid repeated freeze thaw cycles.

#### **APPLICATIONS**

CIB shRNA (m) Lentiviral Particles is recommended for the inhibition of CIB expression in mouse cells.

#### **SUPPORT REAGENTS**

Control shRNA Lentiviral Particles: sc-108080. Available as 200  $\mu$ l frozen viral stock containing 1.0 x 10 $^6$  infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

#### **GENE EXPRESSION MONITORING**

CIB (E-9): sc-271041 is recommended as a control antibody for monitoring of CIB gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **RT-PCR REAGENTS**

Semi-quantitative RT-PCR may be performed to monitor CIB gene expression knockdown using RT-PCR Primer: CIB (m)-PR: sc-43272-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

#### **BIOSAFETY**

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

#### **RESEARCH USE**

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

#### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**