

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

γ2-Syntrophin siRNA (m): sc-43444



BACKGROUND

The syntrophins are structurally related PDZ-domain-containing proteins that facilitate the recruitment of signaling proteins, such as NOS1, to the cell membrane. Syntrophins associate directly with dystrophin, a scaffold protein that is part of a complex which is disrupted in muscular dystrophy, and with dystrophin-related proteins. γ 2-Syntrophin, also known as SYN5 or G2SYN, is a 539 amino acid member of the syntrophin family that functions as an adaptor protein to link and organize various proteins, such as dystrophin, within the cell. γ 2-Syntrophin, which is localized to the sarcolemma cell membrane, is widely expressed and contains one PH domain and one PDZ domain. γ 2-Syntrophin is able to recruit protein to the membrane through its PDZ domain, which is unavailable when the protein is bound to a substrate. As a result of its interaction with various proteins, γ 2-Syntrophin is implicated in inherited muscular dystrophy and in the development of autism.

REFERENCES

- Newey, S.E., Benson, M.A., Ponting, C.P., Davies, K.E. and Blake, D.J. 2000. Alternative splicing of dystrobrevin regulates the stoichiometry of syntrophin binding to the dystrophin protein complex. Curr. Biol. 10: 1295-1298.
- Abdelmoity, A., Padre, R.C., Burzynski, K.E., Stull, J.T. and Lau, K.S. 2000. Neuronal nitric oxide synthase localizes through multiple structural motifs to the sarcolemma in mouse myotubes. FEBS Lett. 482: 65-70.
- 3. Adams, M.E., Kramarcy, N., Krall, S.P., Rossi, S.G., Rotundo, R.L., Sealock, R. and Froehner, S.C. 2000. Absence of α -Syntrophin leads to structurally aberrant neuromuscular synapses deficient in utrophin. J. Cell Biol. 150: 1385-1398.
- Ort, T., Maksimova, E., Dirkx, R., Kachinsky, A.M., Berghs, S., Froehner, S.C. and Solimena, M. 2000. The receptor tyrosine phosphatase-like protein ICA512 binds the PDZ domains of β2-Syntrophin and nNOS in pancreatic β cells. Eur. J. Cell Biol. 79: 621-630.
- Rocco, P., Vainzof, M., Froehner, S.C., Peters, M.F., Marie, S.K., Passos-Bueno, M.R. and Zatz, M. 2000. Brazilian family with pure autosomal dominant spastic paraplegia maps to 8q: analysis of muscle β1-Syntrophin. Am. J. Med. Genet. 92: 122-127.

CHROMOSOMAL LOCATION

Genetic locus: Sntg2 (mouse) mapping to 12 B1.

PRODUCT

 $\gamma 2$ -Syntrophin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see $\gamma 2$ -Syntrophin shRNA Plasmid (m): sc-43444-SH and $\gamma 2$ -Syntrophin shRNA (m) Lentiviral Particles: sc-43444-V as alternate gene silencing products.

For independent verification of γ 2-Syntrophin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43444A, sc-43444B and sc-43444C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

 γ 2-Syntrophin siRNA (m) is recommended for the inhibition of γ 2-Syntrophin expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor γ 2-Syntrophin gene expression knockdown using RT-PCR Primer: γ 2-Syntrophin (m)-PR: sc-43444-PR (20 µl, 440 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.