



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CAPON siRNA (h): sc-43660

BACKGROUND

CAPON (carboxy-terminal PDZ ligand of nNOS) selectively binds within the 100 amino acid PDZ domain of the neuronal nitric oxide synthase (nNOS), but not to endothelial NOS or inducible NOS, and sequesters nNOS in the cytosol. Biosynthesis of the neurotransmitter nitric oxide (NO) requires the association of nNOS with various synaptic proteins, including syntrophin, postsynaptic density (PSD)-95 and PSD-93 through a scaffolding PDZ domain. These proteins facilitate the transport of nNOS to the plasma membrane, where it is catalytically activated by NMDA-receptor mediated calcium channels. The association of nNOS with PSD-95 or PSD-93 is regulated by CAPON. The carboxy-terminus of CAPON binds to the PDZ domain, competes with PSD-95 and PSD-93 for binding to nNOS and in turn prevents the translocation and catalytic activation of nNOS.

REFERENCES

1. Kornau, H.C., Schenker, L.T., Kennedy, M.B. and Seeburg P.H. 1995. Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95. *Science* 269: 1737-1740.
2. Stricker, N.L., Christopherson, K.S., Yi, B.A., Schatz, P.J., Raab, R.W., Dawes, G., Basset, D.E., Jr., Bredt, D.S. and Li, M. 1997. PDZ domain of neuronal nitric oxide synthase recognizes novel C-terminal peptide sequences. *Nat. Biotechnol.* 15: 336-342.
3. Jaffrey, S.R., Snowman, A.M., Eliasson, M.J., Cohen, N.A. and Snyder, S.H. 1998. CAPON: a protein associated with neuronal nitric oxide synthase that regulates its interactions with PSD95. *Neuron* 20: 115-124.
4. Hashida-Okumura, A., Okumura, N., Iwamatsu, A., Buijs, R.M., Romijn, H.J. and Nagai, K. 1999. Interaction of neuronal nitric-oxide synthase with α 1-syntrophin in rat brain. *J. Biol. Chem.* 274: 11736-11741.
5. Oschkinat, H. 1999. A new type of PDZ domain recognition. *Nat. Struct. Biol.* 6: 408-410.

CHROMOSOMAL LOCATION

Genetic locus: NOS1AP (human) mapping to 1q23.3.

PRODUCT

CAPON siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CAPON shRNA Plasmid (h): sc-43660-SH and CAPON shRNA (h) Lentiviral Particles: sc-43660-V as alternate gene silencing products.

For independent verification of CAPON (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43660A, sc-43660B and sc-43660C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CAPON siRNA (h) is recommended for the inhibition of CAPON expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

CAPON (C-9): sc-374504 is recommended as a control antibody for monitoring of CAPON gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CAPON gene expression knockdown using RT-PCR Primer: CAPON (h)-PR: sc-43660-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.