



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



CD52 siRNA (h): sc-44666

BACKGROUND

CD52 is a glycosylphosphatidylinositol (GPI)-linked surface antigen present at high levels on epithelial cells lining the male reproductive tract, thymocytes, lymphocytes, monocytes and macrophages. It is also present at variable levels on lymphoid malignancies. During sperm maturation, spermatozoa passing through the genital tract acquire CD52 that is shed from the epithelial cell lining into seminal plasma. CD52 is detectable on the surface of epididymal sperm and in the ejaculate but not on spermatogenetic cells or testicular spermatozoa. The peptide backbone of CD52, which consists of 12 amino acids, is considered a mere scaffold for posttranslational modifications, such as GPI-anchor and N-glycosylation.

REFERENCES

1. Yeung, C.H., Cooper, T.G. and Nieschlag, E. 1997. Human epididymal secreted protein CD52 on ejaculated spermatozoa: correlations with semen characteristics and the effect of its antibody. *Mol. Hum. Reprod.* 3: 1045-1051.
2. Domagala, A. and Kurpisz, M. 2001. CD52 antigen—a review. *Med. Sci. Monit.* 7: 325-331.
3. Shao, N., Xue, J. and Guo, Z. 2003. Chemical synthesis of CD52 glycopeptides containing the acid-labile fucosyl linkage. *J. Org. Chem.* 68: 9003-9011.
4. Kumar S, Kimlinger, T.K., Lust, J.A., Donovan, K. and Witzig, T.E. 2003. Expression of CD52 on plasma cells in plasma cell proliferative disorders. *Blood* 102: 1075-1077.
5. Hasegawa, A., Sawai, H., Tsubamoto, H., Hori, M., Isojima, S. and Koyama, K. 2004. Possible presence of O-linked carbohydrate in the human male reproductive tract CD52. *J. Reprod. Immunol.* 62: 91-100.

CHROMOSOMAL LOCATION

Genetic locus: CD52 (human) mapping to 1p36.11.

PRODUCT

CD52 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CD52 shRNA Plasmid (h): sc-44666-SH and CD52 shRNA (h) Lentiviral Particles: sc-44666-V as alternate gene silencing products.

For independent verification of CD52 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44666A, sc-44666B and sc-44666C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CD52 siRNA (h) is recommended for the inhibition of CD52 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

CD52 (HI186): sc-51560 is recommended as a control antibody for monitoring of CD52 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG_k BP-HRP: sc-516102 or m-IgG_k BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG_k BP-FITC: sc-516140 or m-IgG_k BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CD52 gene expression knockdown using RT-PCR Primer: CD52 (h)-PR: sc-44666-PR (20 µl, 356 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.