

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



IκB-ζ siRNA (h): sc-44896



The Power to Question

BACKGROUND

IκB- ξ (also called MAIL-S or INAP) is a member of the IκB family. It shares a 30% identity with other family members and consists of six ankyrin repeats at its C-terminal. IκB- ξ accumulates in the nucleus and, in humans, associates with the p50 and p65 subunits of nuclear NFκB via its ankyrin repeats. The mouse homologue of IκB- ξ has only been shown to associate with the p50 subunit. IκB- ξ inhibits DNA binding and activity of the transcription factor NFκB. Distinct from other IκB family members, IκB- ξ is not degraded upon cell stimulation and activation of NFκB, rather evidence shows that it is upregulated under these circumstances. This suggests that IκB- ξ plays a significant role in regulation of NFκB and that NFκB may regulate IκB- ξ in a negative feedback loop. Regulation of NFκB by IκB- ξ may differ depending on the species.

REFERENCES

- 1. Yamazaki, S., et al. 2001. A novel I_KB protein, I_KB - ζ , induced by proinflammatory stimuli, negatively regulates NF $_KB$ in the nuclei. J. Biol. Chem. 276: 27657-27662.
- 2. Muta, T., et al. 2003. I_KB - ξ , a new anti-inflammatory nuclear protein induced by lipopolysaccharide, is a negative regulator for NF $_K$ B. J. Endotoxin. Res. 9: 187-191.
- Shiina, T., et al. 2004. Targeted disruption of MAIL, a nuclear IκB protein, leads to severe atopic dermatitis-like disease. J. Biol. Chem. 279: 55493-55498
- 4. Kusaka, M., et al. 2005. Gene expression profile in rat renal isografts from brain dead donors. Transplant. Proc. 37: 364-366.
- Yamazaki, S., et al. 2005. Stimulus-specific induction of a novel nuclear factor-κB regulator, IκB-ζ, via Toll/Interleukin-1 receptor is mediated by mRNA stabilization. J. Biol. Chem. 280: 1678-1687.
- 6. Motoyama, M., et al. 2005. Positive and negative regulation of nuclear factor- κ B-mediated transcription by $I\kappa$ B- ζ , an inducible nuclear protein. J. Biol. Chem. 280: 7444-7451.
- 7. Muta, T., et al. 2006. $I\kappa B$ - ξ : an inducible regulator of nuclear factor- κB . Vitam. Horm. 74: 301-316.
- 8. Cowland, J.B., et al. 2006. IL-1 β -specific up-regulation of neutrophil gelatinase-associated lipocalin is controlled by I κ B- ζ . J. Immunol. 176: 5559-5566.
- Totzke, G., et al. 2006. A novel member of the lκB family, human lκB-ζ, inhibits transactivation of p65 and its DNA binding. J. Biol. Chem. 281: 12645-12654.

CHROMOSOMAL LOCATION

Genetic locus: NFKBIZ (human) mapping to 3q12.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

IκB- ξ siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IκB- ξ shRNA Plasmid (h): sc-44896-SH and IκB- ξ shRNA (h) Lentiviral Particles: sc-44896-V as alternate gene silencing products.

For independent verification of $l\kappa B$ - ζ (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44896A, sc-44896B and sc-44896C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IkB- $\!\zeta$ siRNA (h) is recommended for the inhibition of IkB- $\!\zeta$ expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor l_KB - ζ gene expression knockdown using RT-PCR Primer: l_KB - ζ (h)-PR: sc-44896-PR (20 μ l, 554 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com