



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PEN-2 siRNA (h): sc-45986

BACKGROUND

Four proteins comprise the γ -secretase complex: Presenilin, nicastrin, Aph-1 and PEN-2. Together, these proteins mediate cell surface signaling pathways for a variety of type I membrane proteins, notably amyloid- β precursor protein, a protein implicated in the development of Alzheimer's disease, via intramembrane proteolysis. The proteins assemble into a proteolytically active complex in the Golgi/*trans*-Golgi network (TGN) compartments. Assembly leads to autocleavage of Presenilin into two subunits to create the active site of γ -secretase, an important step in understanding the mechanisms involved in the etiology and possible treatment of Alzheimer's disease.

REFERENCES

- Kimberly, W.T., et al. 2003. Identity and function of γ -secretase. *J. Neurosci. Res.* 74: 353-360.
- Baulac, S., et al. 2003. Functional γ -secretase complex assembly in Golgi/*trans*-Golgi network: interactions among Presenilin, nicastrin, Aph-1, PEN-2 and γ -secretase substrates. *Neurobiol. Dis.* 14: 194-204.
- Wolfe, M.S. 2003. γ -secretase — intramembrane protease with a complex. *Sci. Aging Knowledge Environ.* 11: PE7.
- Fortna, R.R., et al. 2004. Membrane topology and nicastrin-enhanced endoproteolysis of Aph-1, a component of the γ -secretase complex. *J. Biol. Chem.* 279: 3685-3693.
- Bergman, A., et al. 2004. PEN-2 is sequestered in the endoplasmic reticulum and subjected to ubiquitylation and proteasome-mediated degradation in the absence of Presenilin. *J. Biol. Chem.* 279: 16744-16753.
- Prokop, S., et al. 2004. Requirement of PEN-2 for stabilization of the Presenilin N/C terminal fragment heterodimer within the γ -secretase complex. *J. Biol. Chem.* 279: 23255-23261.
- Hasegawa, H., et al. 2004. Both the sequence and length of the C terminus of PEN-2 are critical for intermolecular interactions and function of Presenilin complexes. *J. Biol. Chem.* 279: 46455-46463.
- Shiraishi, H., et al. 2004. PEN-2 enhances γ -cleavage after Presenilin heterodimer formation. *J. Neurochem.* 90: 1402-1413.
- Kim, S.H. 2005. A sequence within the first transmembrane domain of PEN-2 is critical for PEN-2-mediated endoproteolysis of Presenilin 1. *J. Biol. Chem.* 280: 1992-2001.

CHROMOSOMAL LOCATION

Genetic locus: PSENEN (human) mapping to 19q13.12.

PRODUCT

PEN-2 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PEN-2 shRNA Plasmid (h): sc-45986-SH and PEN-2 shRNA (h) Lentiviral Particles: sc-45986-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PEN-2 siRNA (h) is recommended for the inhibition of PEN-2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

PEN-2 (1C12-G5): sc-293392 is recommended as a control antibody for monitoring of PEN-2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PEN-2 gene expression knockdown using RT-PCR Primer: PEN-2 (h)-PR: sc-45986-PR (20 μ l, 411 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.