



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MCP-1 siRNA (r): sc-45994

BACKGROUND

Eotaxin and the monocyte chemotactic proteins, MCP-1–5, form a subfamily of the C-C (or β) chemokines, which are characterized by a set of conserved adjacent cysteines. MCPs are produced by a variety of cells, including T lymphocytes, subsequent to their activation with cytokines such as IL-1, TNF α and IFN- γ . *In vitro* studies have shown that the MCP isoforms exhibit their chemotactic effects on different subpopulations of lymphocytes. MCP-1 is a potent basophil activator but does not affect eosinophils. MCP-1 levels are increased during infection and inflammation, which are both characterized by leukocyte infiltration. Two MCP-1 receptors, which differ in their carboxy-termini, have been identified.

REFERENCES

1. Charo, I.F., et al. 1994. Molecular cloning and functional expression of two monocyte chemoattractant protein -1 receptors reveals alternative splicing of the carboxyl-terminal tails. *Proc. Natl. Acad. Sci. USA* 91: 2752-2756.
2. Taub, D.D., et al. 1995. Monocyte chemotactic protein-1 (MCP-1), -2 and -3 are chemotactic for human T lymphocytes. *J. Clin. Invest.* 95: 1370-1376.
3. Weber, M., et al. 1995. Monocyte chemotactic protein MCP-2 activates human basophil and eosinophil leukocytes similar to MCP-3. *J. Immunol.* 154: 4166-4172.
4. Combadiere, C., et al. 1995. Monocyte chemoattractant protein-3 is a functional ligand for C-C chemokine receptors 1 and 2B. *J. Biol. Chem.* 270: 29671-29675.
5. Proost, P., et al. 1996. Human monocyte chemotactic proteins-2 and -3: structural and functional comparison with MCP-1. *J. Leukoc. Biol.* 59: 67-74.
6. Dubois, P.M., et al. 1996. Early signal transduction by the receptor to the chemokine monocyte chemotactic protein-1 in a murine T cell hybrid. *J. Immunol.* 156: 1356-1361.
7. Beall, C.J., et al. 1996. Site-directed mutagenesis of monocyte chemoattractant protein-1 identifies two regions of the polypeptide essential for biological activity. *Biochem. J.* 313: 633-640.
8. Kuna, P., et al. 1996. Chemokines in seasonal allergic rhinitis. *J. Allergy Clin. Immunol.* 97: 104-112.

CHROMOSOMAL LOCATION

Genetic locus: Ccl2 (rat) mapping to 10q26.

PRODUCT

MCP-1 siRNA (r) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MCP-1 shRNA Plasmid (r): sc-45994-SH and MCP-1 shRNA (r) Lentiviral Particles: sc-45994-V as alternate gene silencing products.

For independent verification of MCP-1 (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45994A, sc-45994B and sc-45994C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MCP-1 siRNA (r) is recommended for the inhibition of MCP-1 expression in rat cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MCP-1 gene expression knockdown using RT-PCR Primer: MCP-1 (r)-PR: sc-45994-PR (20 μ l, 523 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Zhang, F., et al. 2009. Transforming growth factor- β promotes recruitment of bone marrow cells and bone marrow-derived mesenchymal stem cells through stimulation of MCP-1 production in vascular smooth muscle cells. *J. Biol. Chem.* 284: 17564-17574.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.