



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



HSPB2 (F-9): sc-514154

BACKGROUND

The heat shock proteins (HSPs) comprise a group of highly conserved, abundantly expressed proteins with diverse functions, including the assembly and sequestering of multi-protein complexes, transportation of nascent polypeptide chains across cellular membranes and regulation of protein folding. HSPB2 (heat shock 27 kDa protein 2), also known as HSP 27 or MKBP, is a 182 amino acid protein that belongs to the heat shock protein family and is expressed preferentially in heart and skeletal muscle. Localized to mitochondria, HSPB2 functions as an ATP-dependent chaperone protein that plays a role in the refolding of denatured proteins and may also interact with the Actin cytoskeleton and prevent apoptotic cell death. HSPB2 is abundantly expressed in several cancer cell lines, suggesting that HSPB2 may be an important factor in tumor transformation and metastasis.

REFERENCES

1. Suzuki, A., et al. 1998. MKBP, a novel member of the small heat shock protein family, binds and activates the myotonic dystrophy protein kinase. *J. Cell Biol.* 140: 1113-1124.
2. Concannon, C.G., et al. 2003. On the role of HSP27 in regulating apoptosis. *Apoptosis* 8: 61-70.
3. Wen, F.C., et al. 2003. Down-regulation of heat shock protein 27 in neuronal cells and non-neuronal cells expressing mutant ataxin-3. *FEBS Lett.* 546: 307-314.
4. Mao, H., et al. 2003. HSP72 inhibits focal adhesion kinase degradation in ATP-depleted renal epithelial cells. *J. Biol. Chem.* 278: 18214-18220.
5. Parcellier, A., et al. 2003. HSP27 is a ubiquitin-binding protein involved in IκB-α proteasomal degradation. *Mol. Cell. Biol.* 23: 5790-5802.

CHROMOSOMAL LOCATION

Genetic locus: HSPB2 (human) mapping to 11q23.1; Hspb2 (mouse) mapping to 9 A5.3.

SOURCE

HSPB2 (F-9) is a mouse monoclonal antibody raised against amino acids 36-110 mapping within an internal region of HSPB2 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

HSPB2 (F-9) is available conjugated to agarose (sc-514154 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-514154 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-514154 PE), fluorescein (sc-514154 FITC), Alexa Fluor® 488 (sc-514154 AF488), Alexa Fluor® 546 (sc-514154 AF546), Alexa Fluor® 594 (sc-514154 AF594) or Alexa Fluor® 647 (sc-514154 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-514154 AF680) or Alexa Fluor® 790 (sc-514154 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

HSPB2 (F-9) is recommended for detection of HSPB2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HSPB2 siRNA (h): sc-96438, HSPB2 siRNA (m): sc-146101, HSPB2 shRNA Plasmid (h): sc-96438-SH, HSPB2 shRNA Plasmid (m): sc-146101-SH, HSPB2 shRNA (h) Lentiviral Particles: sc-96438-V and HSPB2 shRNA (m) Lentiviral Particles: sc-146101-V.

Molecular Weight (predicted) of HSPB2: 20 kDa.

Molecular Weight (observed) of HSPB2: 24 kDa.


Positive Controls: human skeletal muscle extract: sc-363776 or human heart extract: sc-363763.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG_κ BP-HRP: sc-516102 or m-IgG_κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG_κ BP-FITC: sc-516140 or m-IgG_κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



HSPB2 (F-9): sc-514154. Western blot analysis of HSPB2 expression in human skeletal muscle (**A**) and human heart (**B**) tissue extracts.



HSPB2 (F-9): sc-514154. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic and nuclear localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.