



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IL-33 (Nessy-1): sc-517600

BACKGROUND

The interleukins (ILs) are a broad family of well characterized cytokines, primarily of hematopoietic cell origin. They are secreted by immune cells (mainly macrophages, B-cells or T-cells) that regulate a wide range of immune system functions. The specific functions of different interleukins vary from the regulation of inflammatory and immune responses to the regulation of other interleukins. IL-33 (interleukin 33), also known as DVS27, IL1F11, C9orf26 or NF-HEV, is a 270 amino acid secreted protein that belongs to the IL family. Expressed in tonsils and lymph nodes, IL-33 functions as a cytokine that is stimulated via interaction with ST2, an association that recruits a variety of proteins, including MyD88, IRAK-1 and TRAF6, and can also induce T helper-associated cytokine activity. IL-33 plays a role in the survival and adhesion of mast cells and may be involved in the control of endothelial cell activation.

REFERENCES

- Baekkeveld, E.S., Roussigne, M., Yamanaka, T., Johansen, F.E., Jahnsen, F.L., Amalric, F., Brandtzaeg, P., Erard, M., Haraldsen, G. and Girard, J.P. 2003. Molecular characterization of NF-HEV, a nuclear factor preferentially expressed in human high endothelial venules. *Am. J. Pathol.* 163: 69-79.
- Hayakawa, H., Hayakawa, M., Kume, A. and Tominaga, S. 2007. Soluble ST2 blocks interleukin-33 signaling in allergic airway inflammation. *J. Biol. Chem.* 282: 26369-26380.
- Sanada, S., Hakuno, D., Higgins, L.J., Schreiter, E.R., McKenzie, A.N. and Lee, R.T. 2007. IL-33 and ST2 comprise a critical biomechanically induced and cardioprotective signaling system. *J. Clin. Invest.* 117: 1538-1549.
- Allakhverdi, Z., Smith, D.E., Comeau, M.R. and Delespesse, G. 2007. Cutting edge: The ST2 ligand IL-33 potently activates and drives maturation of human mast cells. *J. Immunol.* 179: 2051-2054.
- Iikura, M., Suto, H., Kajiwara, N., Oboki, K., Ohno, T., Okayama, Y., Saito, H., Galli, S.J. and Nakae, S. 2007. IL-33 can promote survival, adhesion and cytokine production in human mast cells. *Lab. Invest.* 87: 971-978.
- Küchler, A.M., Pollheimer, J., Balogh, J., Sponheim, J., Manley, L., Sorensen, D.R., De Angelis, P.M., Scott, H. and Haraldsen, G. 2008. Nuclear interleukin-33 is generally expressed in resting endothelium but rapidly lost upon angiogenic or proinflammatory activation. *Am. J. Pathol.* 173: 1229-1242.
- Palmer, G., Lipsky, B.P., Smithgall, M.D., Meininger, D., Siu, S., Talbot-Ayer, D., Gabay, C. and Smith, D.E. 2008. The IL-1 receptor accessory protein (AcP) is required for IL-33 signaling and soluble AcP enhances the ability of soluble ST2 to inhibit IL-33. *Cytokine* 42: 358-364.
- Cherry, W.B., Yoon, J., Bartemes, K.R., Iijima, K. and Kita, H. 2008. A novel IL-1 family cytokine, IL-33, potently activates human eosinophils. *J. Allergy Clin. Immunol.* 121: 1484-1490.
- Kakkar, R. and Lee, R.T. 2008. The IL-33/ST2 pathway: therapeutic target and novel biomarker. *Nat. Rev. Drug Discov.* 7: 827-840.

CHROMOSOMAL LOCATION

Genetic locus: IL33 (human) mapping to 9p24.1; Il33 (mouse) mapping to 19 C1.

SOURCE

IL-33 (Nessy-1) is a mouse monoclonal antibody raised against recombinant IL-33 of human origin.

PRODUCT

Each vial contains 50 µg IgG₁ kappa light chain in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


IL-33 (Nessy-1) is recommended for detection of IL-33 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-33 siRNA (h): sc-75333, IL-33 siRNA (m): sc-75334, IL-33 shRNA Plasmid (h): sc-75333-SH, IL-33 shRNA Plasmid (m): sc-75334-SH, IL-33 shRNA (h) Lentiviral Particles: sc-75333-V and IL-33 shRNA (m) Lentiviral Particles: sc-75334-V.

Molecular Weight of IL-33: 30 kDa.

Positive Controls: human bronchus extract: sc-363754 or human stomach extract: sc-363780.

DATA



IL-33 (Nessy-1): sc-517600. Western blot analysis of IL-33 expression in human bronchus (A) and human stomach (B) tissue extracts.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.