



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# A cyclase III (C-5): sc-518064

## BACKGROUND

Adenylyl cyclases function to convert ATP to cyclic AMP in response to activation by a variety of hormones, neurotransmitters and other regulatory molecules. Cyclic AMP, in turn, activates several other target molecules to control a broad range of diverse phenomena such as metabolism, gene transcription and memory. Adenylyl cyclases respond to receptor-initiated signals, mediated by the G<sub>s</sub> and G<sub>i</sub> heterotrimeric G proteins. The binding of an agonist to a G<sub>s</sub>-coupled receptor catalyzes the exchange of GDP (bound to G<sub>αs</sub>) for GTP, the dissociation of GTP-G<sub>αs</sub> from G<sub>βγ</sub> and G<sub>αs</sub>-mediated activation of adenylyl cyclase. Adenylyl cyclases of the type II family differ from other subforms in that they are conditionally stimulated by G<sub>αs/βγ</sub> subunits and regulated by PKC-mediated C-terminal phosphorylation. Both short- and long-term activation of D<sub>2L</sub> dopamine receptors result in a marked degree of sensitization of A cyclase I, II, V and IX, but not A cyclase VIII. The effects on A cyclase I, II and VIII is dependent upon the ability of these A cyclase isoforms to synergistically respond to selective activators in the presence of activated G<sub>αs</sub>. Belonging to the adenylyl cyclase class IV family, A cyclase III is activated by G<sub>olf</sub>, which results in an elevation of cyclic AMP and subsequent activation of a cyclic nucleotide-gated channel.

## REFERENCES

1. Gilman, A.G. 1987. G proteins: transducers of receptor-generated signals. *Annu. Rev. Biochem.* 56: 615-649.
2. Bourne, H.R., et al. 1990. The GTPase superfamily: a conserved switch for diverse cell functions. *Nature* 348: 125-132.
3. Tang, W.J., et al. 1992. Adenylyl cyclases. *Cell* 70: 869-872.
4. Taussig, R., et al. 1994. Distinct patterns of bidirectional regulation of mammalian adenylyl cyclases. *J. Biol. Chem.* 269: 6093-6100.
5. Liu, C.Y., et al. 1999. FICRHR/cyclic AMP signaling in myenteric ganglia and calbindin-D28 intrinsic primary afferent neurons involves adenylyl cyclases I, III and IV. *Brain Res.* 826: 253-269.
6. Gibson, A.D. and Garbers, D.L. 2000. Guanylyl cyclases as a family of putative odorant receptors. *Annu. Rev. Neurosci.* 23 : 417-439.
7. Parkinson, N.A., et al. 2001. A nuclear location for Ca<sup>2+</sup>-activated adenylyl cyclases I and III in neurons. *Brain Res. Mol. Brain Res.* 91: 43-49.
8. Cumbay, M.G., et al. 2001. Heterologous sensitization of recombinant adenylate cyclases by activation of D<sub>2</sub> dopamine receptors. *J. Pharmacol. Exp. Ther.* 297: 1201-1209.

## CHROMOSOMAL LOCATION

Genetic locus: ADCY3 (human) mapping to 2p23.3.

## SOURCE

A cyclase III (C-5) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 22-46 within an N-terminal cytoplasmic domain of A cyclase III of human origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200 µg IgM in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS


A cyclase III (C-5) is recommended for detection of A cyclase III of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for A cyclase III siRNA (h): sc-29600, A cyclase III shRNA Plasmid (h): sc-29600-SH and A cyclase III shRNA (h) Lentiviral Particles: sc-29600-V.

Molecular Weight of A cyclase III glycosylated forms: 170/180 kDa.

Positive Controls: mouse A cyclase III transfected 293T whole cell lysate, IMR-32 cell lysate: sc-2409 or WI-38 whole cell lysate: sc-364260.

## DATA



A cyclase III (C-5): sc-518064. Western blot analysis of A cyclase III expression in non-transfected 293T (**A**) and mouse A cyclase III transfected 293T (**B**) whole cell lysates.

A cyclase III (C-5): sc-518064. Western blot analysis of A cyclase III expression in non-transfected 293T (**A**) and mouse A cyclase III transfected 293T (**B**) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.