



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



SDF-1 (F-4): sc-518066



The Power to Question

BACKGROUND

The C-X-C or α chemokine family is characterized by a pair of cysteine residues separated by a single amino acid and primarily functions as chemoattractants for neutrophils. The C-X-C family includes IL-8, NAP-2, MSGA and stromal cell-derived factor-1, or SDF-1. SDF-1 was originally described as a pre-B cell stimulatory factor, but has now been shown to function as a potent chemoattractant for T cells and monocytes, but not neutrophils. Receptors for the C-X-C family are G protein-coupled, seven-pass, transmembrane domain proteins which include IL-8RA, IL-8RB and fusin (also designated LESTR or CXCR-4). Fusin is highly homologous to the IL-8 receptors, sharing 37% sequence identity at the amino acid level. The IL-8 receptors bind to IL-8, NAP-2 and MSGA, while fusin binds to its cognate ligand, SDF-1. Fusin has been identified as the major co-receptor for T-tropic HIV-1, and SDF-1 has been shown to inhibit HIV-1 infection. Six human SDF-1 isoforms exist due to alternative splicing of CXCL12, the gene encoding SDF-1. Three isoforms are known for mouse and rat.

REFERENCES

- Laterveer, L., et al. 1996. Rapid mobilization of hematopoietic progenitor cells in rhesus monkeys by a single intravenous injection of Interleukin-8. *Blood* 87: 781-788.
- Deng, H., et al. 1996. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 381: 661-666.
- Nagasaki, T., et al. 1996. Defects of B cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the C-X-C chemokine PBSF/SDF-1. *Nature* 382: 635-638.
- Bleul, C.C., et al. 1996. The lymphocyte chemoattractant SDF-1 is a ligand for LESTR/fusin and blocks HIV-1 entry. *Nature* 382: 829-833.
- Feng, Y., et al. 1996. HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 272: 872-877.

CHROMOSOMAL LOCATION

Genetic locus: CXCL12 (human) mapping to 10q11.21.

SOURCE

SDF-1 (F-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 66-92 at the C-terminus of SDF-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

SDF-1 (F-4) is available conjugated to agarose (sc-518066 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-518066 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-518066 PE), fluorescein (sc-518066 FITC), Alexa Fluor[®] 488 (sc-518066 AF488), Alexa Fluor[®] 546 (sc-518066 AF546), Alexa Fluor[®] 594 (sc-518066 AF594) or Alexa Fluor[®] 647 (sc-518066 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-518066 AF680) or Alexa Fluor[®] 790 (sc-518066 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

SDF-1 (F-4) is recommended for detection of all known isoforms of SDF-1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SDF-1 siRNA (h): sc-39367, SDF-1 shRNA Plasmid (h): sc-39367-SH and SDF-1 shRNA (h) Lentiviral Particles: sc-39367-V.


Molecular Weight of SDF-1: 10 kDa.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgG_x BP-HRP: sc-516102 or m-IgG_x BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™
- Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- Immunofluorescence: use m-IgG_x BP-FITC: sc-516140 or m-IgG_x BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



SDF-1 (F-4): sc-518066. Western blot analysis of SDF-1 expression in human SDF-1 α sc-4654 (**A**) and human SDF-1 β sc-4655 (**B**) fusion proteins. Detection reagent used: m-IgG_x BP-HRP (Cruz Marker): sc-516102-CM.



SDF-1 (F-4): sc-518066. Western blot analysis of human recombinant SDF-1 fusion protein.

SELECT PRODUCT CITATIONS

- Zhou, W., et al. 2020. Discovery of exosomes from tick saliva and salivary glands reveals therapeutic roles for CXCL12 and IL-8 in wound healing at the tick-human skin interface. *Front. Cell Dev. Biol.* 8: 554.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA