



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# GOS2 (D-1): sc-518067



The Power to Question

## BACKGROUND

GOS2 (putative lymphocyte G<sub>0</sub>/G<sub>1</sub> switch protein 2) is a 103 amino acid novel target of peroxisome proliferator-activated receptors (PPARs) and regulator of latent HIV. GOS2 may be involved in adipocyte differentiation and its expression is essential for committing cells to enter the G<sub>1</sub> phase of the cell cycle. GOS2 contains a CpG-rich island and multiple sites for potential phosphorylation by casein kinase II and protein kinase C. The gene encoding GOS2 maps to human chromosome 1, which is the largest human chromosome. Chromosome 1 spans about 260 million base pairs and makes up 8% of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes Lamin A. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma.

## REFERENCES

- Russell, L. and Forsdyke, D.R. 1991. A human putative lymphocyte G<sub>0</sub>/G<sub>1</sub> switch gene containing a CpG-rich island encodes a small basic protein with the potential to be phosphorylated. *DNA Cell Biol.* 10: 581-591.
- Cristillo, A.D., et al. 1997. Cyclosporin A inhibits early mRNA expression of G<sub>0</sub>/G<sub>1</sub> switch gene 2 (GOS2) in cultured human blood mononuclear cells. *DNA Cell Biol.* 16: 1449-1458.
- Eudy, J.D., et al. 1998. Mutation of a gene encoding a protein with extracellular matrix motifs in Usher syndrome type Ila. *Science* 280: 1753-1757.

## CHROMOSOMAL LOCATION

Genetic locus: GOS2 (human) mapping to 1q32.2.

## SOURCE

GOS2 (D-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-19 at the N-terminus of GOS2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

GOS2 (D-1) is available conjugated to agarose (sc-518067 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-518067 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-518067 PE), fluorescein (sc-518067 FITC), Alexa Fluor® 488 (sc-518067 AF488), Alexa Fluor® 546 (sc-518067 AF546), Alexa Fluor® 594 (sc-518067 AF594) or Alexa Fluor® 647 (sc-518067 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-518067 AF680) or Alexa Fluor® 790 (sc-518067 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

GOS2 (D-1) is recommended for detection of GOS2 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GOS2 siRNA (h): sc-78689, GOS2 shRNA Plasmid (h): sc-78689-SH and GOS2 shRNA (h) Lentiviral Particles: sc-78689-V.


Molecular Weight of GOS2: 11 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG<sub>κ</sub> BP-HRP: sc-516102 or m-IgG<sub>κ</sub> BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG<sub>κ</sub> BP-FITC: sc-516140 or m-IgG<sub>κ</sub> BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



GOS2 (D-1): sc-518067. Western blot analysis of human recombinant GOS2 fusion protein.

## SELECT PRODUCT CITATIONS

- Ma, Y., et al. 2019. Activation of G<sub>0</sub>/G<sub>1</sub> switch gene 2 by endoplasmic reticulum stress enhances hepatic steatosis. *Metab. Clin. Exp.* 99: 32-44.
- van Dierendonck, X.A.M.H., et al. 2020. HILPDA uncouples lipid droplet accumulation in adipose tissue macrophages from inflammation and metabolic dysregulation. *Cell Rep.* 30: 1811.e6-1822.e6.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA