



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NFKBIL1 (A-6): sc-518229

BACKGROUND

NFκB, a pleiotropic transcription factor, is present in almost all cell types and is involved in many biological processes including inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NFκB is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. This complex is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. The NFκB inhibitor-like protein 1 (NFKBIL1), also designated IKBL, acts as a negative regulator of NFκB activation. Mutations in the NFKBIL1 gene have been linked to several disorders including type 1 diabetes, rheumatoid arthritis, ulcerative colitis and chronic Chagas cardiomyopathy.

REFERENCES

1. Ruben, S.M., et al. 1992. Functional characterization of the NFκB p65 transcriptional activator and an alternatively spliced derivative. *Mol. Cell Biol.* 12: 444-454.
2. Deloukas, P. and van Loon, A.P. 1993. Genomic organization of the gene encoding the p65 subunit of NFκB: multiple variants of the p65 protein may be generated by alternative splicing. *Hum. Mol. Genet.* 2: 1895-1900.
3. Handel-Fernandez, M.E. and Vincek, V. 1999. Sequence analysis and expression of a mouse homolog of human IkBL gene. *Biochim. Biophys. Acta* 1444: 306-310.
4. de la Concha, E.G., et al. 2000. Susceptibility to severe ulcerative colitis is associated with polymorphism in the central MHC gene IKBL. *Gastroenterology* 119: 1491-1495.
5. Yamashita, T., et al. 2004. IKBL promoter polymorphism is strongly associated with resistance to type 1 diabetes in Japanese. *Tissue Antigens* 63: 223-230.

CHROMOSOMAL LOCATION

Genetic locus: NFKBIL1 (human) mapping to 6p21.33.

SOURCE

NFKBIL1 (A-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 220-246 of NFKBIL1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

NFKBIL1 (A-6) is available conjugated to agarose (sc-518229 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-518229 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-518229 PE), fluorescein (sc-518229 FITC), Alexa Fluor® 488 (sc-518229 AF488), Alexa Fluor® 546 (sc-518229 AF546), Alexa Fluor® 594 (sc-518229 AF594) or Alexa Fluor® 647 (sc-518229 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-518229 AF680) or Alexa Fluor® 790 (sc-518229 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

NFKBIL1 (A-6) is recommended for detection of NFKBIL1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NFKBIL1 siRNA (h): sc-95606, NFKBIL1 shRNA Plasmid (h): sc-95606-SH and NFKBIL1 shRNA (h) Lentiviral Particles: sc-95606-V.


Molecular Weight of NFKBIL1: 43 kDa.

Positive Controls: NFKBIL1 (h): 293T Lysate: sc-369815.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



NFKBIL1 (A-6): sc-518229. Western blot analysis of NFKBIL1 expression in human recombinant NFKBIL1 fusion protein (A), non-transfected: sc-117752 (B) and human NFKBIL1 transfected: sc-369815 (C) 293T whole cell lysates. Detection reagent used: m-IgG₁ BP-HRP: sc-525408.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.