



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# SIRP- $\gamma$ (OX119): sc-53114

## BACKGROUND

SIRPs (signal-regulatory proteins) are a family of transmembrane glycoproteins that were identified by their association with the Src homology 2 domain-containing protein-tyrosine phosphatase SHP-2 in response to Insulin. The SIRP family negatively regulates the PI 3-K pathway, which may diminish EGFR-mediated motility and survival phenotypes that contribute to transformation of certain cell types. SIRP- $\alpha$ 1 is a transmembrane protein which contains an extracellular portion with three immunoglobulin-like structures and a cyto-plasmic region with four potential tyrosine phosphorylation sites. SIRP- $\alpha$ 1 is a substrate for activated receptor tyrosine kinases. In its tyrosine phosphorylated form, SIRP- $\alpha$ 1 binds to SH-PTP2 through SH2 interactions and acts as an SH-PTP2 substrate. SIRP- $\alpha$ 1 has been shown to have negative regulatory effects on cellular responses induced by growth factors, oncogenes and insulin. SIRP- $\beta$ 1 shares extensive sequence homology with SIRP- $\alpha$ 1 in its extracellular portion but lacks the cytoplasmic portion. SIRP- $\gamma$ , originally designated SIRP- $\beta$ 2 (SIRP-B2, CD172g) has unique characteristics from both the  $\alpha$  and  $\beta$  versions. SIRP- $\gamma$  is expressed on the majority of T cells and a proportion of B cells. CD47 associates with SIRP- $\gamma$ , and this interaction signals unidirectionally only.

## REFERENCES

1. Yamauchi, K., et al. 1995. Identification of the major SHPTP2-binding protein that is tyrosine-phosphorylated in response to Insulin. *J. Biol. Chem.* 270: 17716-17722.
2. Fujioka, Y., et al. 1996. A novel membrane glycoprotein, SHPS-1, that binds the SH2-domain-containing tyrosine phosphatase SHP-2 in response to mitogens and cell adhesion. *Mol. Cell. Biol.* 16: 6887-6899.
3. Kharitonkov, A., et al. 1997. A family of proteins that inhibit signalling through tyrosine kinase receptors. *Nature* 386: 181-186.
4. Stofega, M.R., et al. 1998. Growth hormone regulation of SIRP and SHP-2 tyrosyl phosphorylation and association. *J. Biol. Chem.* 273: 7112-7117.
5. Wu, C.J., et al. 2000. Inhibition of EGFR-mediated phosphoinositide-3-OH kinase (PI-3 K) signaling and glioblastoma phenotype by signal-regulatory proteins (SIRPs). *Oncogene* 19: 3999-4010.
6. Latour, S., et al. 2001. Bidirectional negative regulation of human T and dendritic cells by CD47 and its cognate receptor signal-regulator protein- $\alpha$ : downregulation of IL-12 responsiveness and inhibition of dendritic cell activation. *J. Immunol.* 167: 2547-2554.
7. Brooke, G., et al. 2004. Human lymphocytes interact directly with CD47 through a novel member of the signal regulatory protein (SIRP) family. *J. Immunol.* 173: 2562-2570.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: SIRPG (human) mapping to 20p13.

## SOURCE

SIRP- $\gamma$  (OX119) is a mouse monoclonal antibody raised against recombinant SIRP- $\gamma$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

SIRP- $\gamma$  (OX119) is available conjugated to agarose (sc-53114 AC), 500  $\mu$ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-53114 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-53114 PE), fluorescein (sc-53114 FITC), Alexa Fluor® 488 (sc-53114 AF488), Alexa Fluor® 546 (sc-53114 AF546), Alexa Fluor® 594 (sc-53114 AF594) or Alexa Fluor® 647 (sc-53114 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-53114 AF680) or Alexa Fluor® 790 (sc-53114 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## APPLICATIONS


SIRP- $\gamma$  (OX119) is recommended for detection of SIRP- $\gamma$  of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and flow cytometry (1  $\mu$ g per 1 x 10<sup>6</sup> cells); may cross-react with over-expressed SIRP- $\alpha$  and SIRP- $\beta$ .

Suitable for use as control antibody for SIRP- $\gamma$  siRNA (h): sc-76499, SIRP- $\gamma$  shRNA Plasmid (h): sc-76499-SH and SIRP- $\gamma$  shRNA (h) Lentiviral Particles: sc-76499-V.


Molecular Weight of SIRP- $\gamma$ : 55 kDa.

Positive Controls: SIRP- $\gamma$  (h): 293T Lysate: sc-116900.

## DATA



SIRP- $\gamma$  (OX119): sc-53114. Western blot analysis of SIRP- $\gamma$  expression in non-transfected: sc-117752 (A) and human SIRP- $\gamma$  transfected: sc-116900 (B) 293T whole cell lysates. Detection reagent used: m-IgG<sub>1</sub> BP-HRP: sc-525408.



SIRP- $\gamma$  (OX119): sc-53114. Indirect FCM analysis of human peripheral blood leukocytes stained with SIRP- $\gamma$  (OX119), followed by PE-conjugated goat anti-mouse IgG<sub>1</sub>: sc-3764. Black line histogram represents the isotype control, normal mouse IgG<sub>1</sub>: sc-3877.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.