



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# m-IgG<sub>3</sub> BP-HRP: sc-533670

## BACKGROUND

Mouse IgG<sub>3</sub> binding protein (m-IgG<sub>3</sub> BP) conjugated to horseradish peroxidase (HRP) is a strongly recommended alternative to conventional goat/rabbit anti-mouse IgG secondary antibodies for Western Blotting (WB) and immunohistochemical (IHC) signal enhancement. Mouse IgG<sub>3</sub> binding protein is a highly specific reagent that provides strong signal with minimal background and virtually complete elimination of lot to lot variation associated with conventionally generated secondary antibodies. Mouse IgG<sub>3</sub> binding protein (m-IgG<sub>3</sub> BP) is suitable for binding to most, but not all mouse monoclonal IgG<sub>3</sub> antibodies; not suitable for use with mouse monoclonal IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>2b</sub>, IgM, IgA or IgE antibodies. Not cross reactive with human or rat IgG antibodies.

## SOURCE

m-IgG<sub>3</sub> BP-HRP is a purified recombinant mouse IgG<sub>3</sub> binding protein conjugated to horseradish peroxidase (HRP).

## PRODUCT

Each vial contains 200 µg mouse IgG<sub>3</sub> binding protein-HRP in 0.5 ml of PBS containing 40% glycerol, 1% stabilizer protein and < 0.01% thimerosal.

## APPLICATIONS

m-IgG<sub>3</sub> BP-HRP is recommended for detection of mouse IgG<sub>3</sub> by ECL Western Blotting (starting dilution: 1:1000, dilution range: 1:1000-1:10000) and immunohistochemistry (including paraffin-embedded sections) (starting dilution: 1:25, dilution range 1:25-1:100). Optimal dilution to be determined by titration.

For Western Blotting using tissue extracts and m-IgG<sub>3</sub> BP-HRP, we strongly recommend subtracting endogenous immunoglobulins from extracts with Protein G PLUS-Agarose Reagent: sc-2002, to prevent Western Blotting interference when detecting proteins of approximately 25 kDa in size.

## RECOMMENDED SUPPORT PRODUCTS

- Protein G PLUS-Agarose Reagent: sc-2002
- Western Blotting Luminol Reagent, for 2,000 cm<sup>2</sup> membrane area: sc-2048
- RIPA Lysis Buffer, 50 ml, cell lysis buffer with protease inhibitors: sc-24948
- Electrophoresis Sample Buffer, 2X, 25 ml, reducing buffer: sc-24945
- UltraCruz<sup>®</sup> Protease Inhibitor Cocktail Tablet, 20 tablets: sc-29130
- Running Buffer, 10X, 1 L, TRIS-Glycine WB running buffer, pH 8.3: sc-24949
- Towbin, with SDS, 10X, 1 L, WB transfer buffer pH 8.3: sc-24954
- TBS Blotto A, lyophilized powder in single-use bottle: sc-2333
- UltraCruz<sup>®</sup> PVDF Transfer Membrane, 0.45 µm, 30 cm x 3 m roll: sc-3723
- UltraCruz<sup>®</sup> Nitrocellulose Pure Transfer Membrane, 0.22 µm, 30 cm x 3 m roll: sc-3718
- UltraCruz<sup>®</sup> Tissue Culture Dish, 100 mm polystyrene dish: sc-200286
- UltraCruz<sup>®</sup> Autoradiography Film, Blue, 8 x 1, 100 sheets: sc-201697
- UltraCruz<sup>®</sup> Gel Incubation Trays, 100 per pack: sc-201755 (blue), sc-201756 (green), sc-201757 (pink), sc-201758 (yellow), sc-201759 (orange)

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.