



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

m-IgG₃ BP-CFL 647: sc-533676

BACKGROUND

Mouse IgG₃ binding protein (m-IgG₃ BP) conjugated to CruzFluor™ 647 (CFL 647) is a strongly recommended alternative to conventional goat/rabbit anti-mouse IgG secondary antibodies for RGB Western Blotting (WB), immunofluorescence (IF) and flow cytometry (FCM) signal enhancement. CruzFluor™ 647 (CFL 647) is a far-red fluorescent dye that is an excellent substitute for AlexaFluor® 647, offering comparable photostability and the ability to resist protein quenching. Suitable for use with RGB imaging systems, such as Invitrogen/iBright and other comparable systems. Mouse IgG₃ binding protein is a highly specific reagent that provides strong signal with minimal background and virtually complete elimination of lot to lot variation associated with conventionally generated secondary antibodies. Mouse IgG₃ binding protein (m-IgG₃ BP) is suitable for binding to most, but not all mouse monoclonal IgG₃ antibodies; not suitable for use with mouse monoclonal IgG₁, IgG_{2a}, IgG_{2b}, IgM, IgA or IgE antibodies. Not cross reactive with human or rat IgG antibodies.

SOURCE

m-IgG₃ BP-CFL 647 is a purified recombinant mouse IgG₃ binding protein conjugated to CruzFluor™ 647 (CFL 647).

PRODUCT

Each vial contains 200 µg mouse IgG₃ binding protein-CFL 647 in 0.5 ml of PBS containing 0.1% gelatin and 0.1% sodium azide.

APPLICATIONS

m-IgG₃ BP-CFL 647 is recommended for detection of mouse IgG₃ by RGB Western Blotting (starting dilution: 1:1000, dilution range: 1:500-1:2000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:200) and flow cytometry (0.5-1 µg per 1 x 10⁶ cells). Optimal dilution to be determined by titration.

RECOMMENDED SUPPORT PRODUCTS

- CrystalCruz® Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336
- FCM Lysing solution: sc-3621
- FCM Fixation Buffer: sc-3622
- FCM Permeabilization Buffer: sc-3623
- FCM Wash Buffer: sc-3624
- Intracellular FCM System: sc-45063

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

CRUZFLUOR™ SPECTRAL PROPERTIES

PRODUCT	CAT. #	EXCITATION MAXIMUM	EMISSION MAXIMUM
m-IgG ₃ BP-CFL 488	sc-533673	488 nm	514 nm
m-IgG ₃ BP-CFL 555	sc-533674	556 nm	569 nm
m-IgG ₃ BP-CFL 594	sc-533675	587 nm	603 nm
m-IgG ₃ BP-CFL 647	sc-533676	654 nm	669 nm
m-IgG ₃ BP-CFL 680	sc-533677	683 nm	700 nm
m-IgG ₃ BP-CFL 790	sc-533678	786 nm	811 nm