



SZABO  
SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## m-IgG<sub>2b</sub> BP-FITC: sc-542743

### BACKGROUND

Mouse IgG<sub>2b</sub> binding protein (m-IgG<sub>2b</sub> BP) conjugated to fluorescein isothiocyanate (FITC) is a strongly recommended alternative to conventional goat/rabbit anti-mouse IgG secondary antibodies for immunofluorescence (IF) and flow cytometry (FCM) signal enhancement. Mouse IgG<sub>2b</sub> binding protein is a highly specific reagent that provides strong signal with minimal background and virtually complete elimination of lot to lot variation associated with conventionally generated secondary antibodies. Mouse IgG<sub>2b</sub> binding protein (m-IgG<sub>2b</sub> BP) is suitable for binding to most, but not all mouse monoclonal IgG<sub>2b</sub> antibodies; not suitable for use with mouse monoclonal IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>3</sub>, IgM, IgA or IgE antibodies. Not cross reactive with human, rat or goat IgG antibodies.

### SOURCE

m-IgG<sub>2b</sub> BP-FITC is a purified recombinant mouse IgG<sub>2b</sub> binding protein conjugated to fluorescein isothiocyanate (FITC).

### PRODUCT

Each vial contains 50 µg mouse IgG<sub>2b</sub> binding protein-FITC in 0.5 ml of PBS containing 0.1% gelatin and 0.1% sodium azide.

### APPLICATIONS

m-IgG<sub>2b</sub> BP-FITC is recommended for detection of mouse IgG<sub>2b</sub> by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:200) and flow cytometry (0.5-1 µg per 1 x 10<sup>6</sup> cells). Optimal dilution to be determined by titration.

FITC excitation/emission spectrum peak wavelengths: 495 nm/519 nm

### RECOMMENDED SUPPORT PRODUCTS

- CrystalCruz<sup>®</sup> Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336
- FCM Lysing solution: sc-3621
- FCM Fixation Buffer: sc-3622
- FCM Permeabilization Buffer: sc-3623
- FCM Wash Buffer: sc-3624
- Intracellular FCM System: sc-45063

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.