



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



brachyury 2 siRNA (m): sc-141735

BACKGROUND

The T-box gene family consists of members that share a unique DNA binding domain. The best characterized T-box (Tbx) gene, Brachyury or T, encodes a transcription factor that plays an important role in early vertebrate development. Tbx genes are a family of developmental regulators with more than 20 members recently identified among invertebrates and vertebrates. Mutations in Tbx genes have been found to cause several human diseases. The understanding of functional mechanisms of Tbx products has come mainly from the prototypical T/Brachyury protein, which is a transcription activator. The T-domain is a highly conserved DNA-binding motif originally defined in Brachyury and characteristic of the Tbx family of transcription factors. The murine Brachyury (T) gene is required in posterior mesoderm formation and axial development. Mutant embryos lacking T gene function are deficient in notochord differentiation and posterior mesoderm formation, but develop anterior mesoderm.

REFERENCES

1. Kispert, A. and Herrmann, B.G. 1994. Immunohistochemical analysis of the Brachyury protein in wild-type and mutant mouse embryos. *Dev. Biol.* 161: 179-193.
2. Conlon, F.L., Wright, C.V. and Robertson, E.J. 1995. Effects of the TWist mutation on notochord formation and mesodermal patterning. *Mech. Dev.* 49: 201-209.
3. Agulnik, S.I., Ruvinsky, I. and Silver, L.M. 1997. Three novel T-box genes in *Caenorhabditis elegans*. *Genome* 40: 458-464.
4. He, M.I., Wen, L., Campbell, C.E., Wu, J.Y. and Rao, Y. 1999. Transcription repression by *Xenopus* ET and its human ortholog TBX3, a gene involved in ulnar-mammary syndrome. *Proc. Natl. Acad. Sci. USA* 96: 10212-10217.
5. Hurlin, P.J., Steingrimsson, E., Copeland, N.G., Jenkins, N.A. and Eisenman, R.N. 1999. Mga, a dual-specificity transcription factor that interacts with Max and contains a T-domain DNA-binding motif. *EMBO J.* 18: 7019-7028.

CHROMOSOMAL LOCATION

Genetic locus: T2 (mouse) mapping to 17 A1.

PRODUCT

brachyury 2 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see brachyury 2 shRNA Plasmid (m): sc-141735-SH and brachyury 2 shRNA (m) Lentiviral Particles: sc-141735-V as alternate gene silencing products.

For independent verification of brachyury 2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141735A and sc-141735B.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

brachyury 2 siRNA (m) is recommended for the inhibition of brachyury 2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor brachyury 2 gene expression knockdown using RT-PCR Primer: brachyury 2 (m)-PR: sc-141735-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.