



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

cadherin-18 siRNA (m): sc-141971

BACKGROUND

The cadherins are a family of Ca^{2+} -dependent adhesion molecules that function to mediate cell-cell binding events that are critical to the maintenance of cell structure and morphogenesis. EY-cadherin, also known as CDH18 (cadherin 18), CDH14 (cadherin 14), CDH24 or CDH14L, is a 790 amino acid single-pass type I membrane protein that contains five cadherin domains. One of several members of the cadherin superfamily, EY-cadherin functions as a type II classical cadherin that is expressed specifically in the central nervous system (CNS), where it plays a role in cell-cell binding events. Specifically, EY-cadherin is thought to be involved in axon guidance and outgrowth, as well as synaptic adhesion within the CNS. EY-cadherin contains a highly conserved C-terminal domain characteristic of all cadherins, but lacks the HAV cell adhesion sequence that is specific to type I cadherins. The gene encoding EY-cadherin is located within a region on chromosome five that is commonly deleted in carcinomas, implicating EY-cadherin as a potential tumor suppressor.

REFERENCES

1. Shibata, T., Shimoyama, Y., Gotoh, M. and Hirohashi, S. 1997. Identification of human cadherin-14, a novel neurally specific type II cadherin, by protein interaction cloning. *J. Biol. Chem.* 272: 5236-5240.
2. Kools, P., Vanhalst, K., Van den Eynde, E. and van Roy, F. 1999. The human cadherin-10 gene: complete coding sequence, predominant expression in the brain, and mapping on chromosome 5p13-14. *FEBS Lett.* 452: 328-334.
3. Chalmers, I.J., Höfler, H. and Atkinson, M.J. 1999. Mapping of a cadherin gene cluster to a region of chromosome 5 subject to frequent allelic loss in carcinoma. *Genomics* 57: 160-163.
4. Shimoyama, Y., Takeda, H., Yoshihara, S., Kitajima, M. and Hirohashi, S. 1999. Biochemical characterization and functional analysis of two type II classic cadherins, cadherin-6 and -14, and comparison with E-cadherin. *J. Biol. Chem.* 274: 11987-11994.
5. Shimoyama, Y., Tsujimoto, G., Kitajima, M. and Natori, M. 2000. Identification of three human type-II classic cadherins and frequent heterophilic interactions between different subclasses of type-II classic cadherins. *Biochem. J.* 349: 159-167.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603019. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Aijaz, S., Allen, J., Tregidgo, R., van Heyningen, V., Hanson, I. and Clark, B.J. 2005. Expression analysis of SIX3 and SIX6 in human tissues reveals differences in expression and a novel correlation between the expression of SIX3 and the genes encoding isocitrate dehydrogenase and cadherin 18. *Genomics* 86: 86-99.
8. Heikinheimo, K., Jee, K.J., Morgan, P.R., Nagy, B., Knuutila, S. and Leivo, I. 2007. Genetic changes in sporadic keratocystic odontogenic tumors (odontogenic keratocysts). *J. Dent. Res.* 86: 544-549.

CHROMOSOMAL LOCATION

Genetic locus: Cdh18 (mouse) mapping to 15 A2.

PRODUCT

cadherin-18 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see cadherin-18 shRNA Plasmid (m): sc-141971-SH and cadherin-18 shRNA (m) Lentiviral Particles: sc-141971-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

cadherin-18 siRNA (m) is recommended for the inhibition of cadherin-18 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor cadherin-18 gene expression knockdown using RT-PCR Primer: cadherin-18 (m)-PR: sc-141971-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.