



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



calsyntenin-2 siRNA (m): sc-141986

BACKGROUND

Members of the calsyntenin protein family are localized to the post-synaptic membrane of excitatory central nervous system (CNS) synapses. Calsyntenin-2, also known as Alcadein- γ , is a 955 amino acid protein that localizes to the endoplasmic reticulum, Golgi apparatus and plasma membranes. Containing two cadherin-like repeats in its N-terminal extracellular region, calsyntenin-2 binds synaptic calcium with its cytoplasmic domain, suggesting a role in the modulation of calcium-mediated postsynaptic signals. Under normal physiological conditions, calsyntenin-2 is proteolytically processed in an event in which the primary ζ -cleavage generates a short C-terminal transmembrane fragment and a long extracellular N-terminal domain.

REFERENCES

1. Vogt, L., et al. 2001. Calsyntenin-1, a proteolytically processed postsynaptic membrane protein with a cytoplasmic calcium-binding domain. *Mol. Cell. Neurosci.* 17: 151-166.
2. Hintsch, G., et al. 2002. The calsyntenins—a family of postsynaptic membrane proteins with distinct neuronal expression patterns. *Mol. Cell. Neurosci.* 21: 393-409.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611323. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Araki, Y., et al. 2003. Novel cadherin-related membrane proteins, Alcadeins, enhance the X11-like protein-mediated stabilization of amyloid β -protein precursor metabolism. *J. Biol. Chem.* 278: 49448-49458.
5. Araki, Y., et al. 2004. Coordinated metabolism of Alcadein and amyloid β -protein precursor regulates FE65-dependent gene transactivation. *J. Biol. Chem.* 279: 24343-24354.
6. Suzuki, T., et al. 2006. Trafficking of Alzheimer's disease-related membrane proteins and its participation in disease pathogenesis. *J. Biochem.* 139: 949-955.
7. Cheng, X.R., et al. 2006. The family of calsyntenins: learning and memory related genes. *Sheng Li Ke Xue Jin Zhan* 37: 205-210.

CHROMOSOMAL LOCATION

Genetic locus: Clntn2 (mouse) mapping to 9 E3.3.

PRODUCT

calsyntenin-2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see calsyntenin-2 shRNA Plasmid (m): sc-141986-SH and calsyntenin-2 shRNA (m) Lentiviral Particles: sc-141986-V as alternate gene silencing products.

For independent verification of calsyntenin-2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-141986A, sc-141986B and sc-141986C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

calsyntenin-2 siRNA (m) is recommended for the inhibition of calsyntenin-2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

calsyntenin-2 (T-16): sc-102403 is recommended as a control antibody for monitoring of calsyntenin-2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor calsyntenin-2 gene expression knockdown using RT-PCR Primer: calsyntenin-2 (m)-PR: sc-141986-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.