



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)





CD209c siRNA (m): sc-142186

BACKGROUND

Antigen-presenting cells are localized in essentially every tissue, where they operate at the interface of innate and acquired immunity by capturing pathogens and presenting pathogen-derived peptides to T cells. Dendritic cells capture antigens or viruses in peripheral tissue and transport them to lymphoid organs, an event that induces cellular T cell responses. The mouse CD209 family of cell adhesion receptors consists of CD209a (also known as DC-SIGN), CD209b, CD209c, CD209d, CD209e, CD209f and CD209g, all of which function to mediate the endocytosis and subsequent degradation of pathogens within lysosomal compartments. There are two human CD209 proteins, designated DC-SIGN and DC-SIGNR, which function in a similar manner to the mouse proteins.

REFERENCES

1. Engering, A., et al. 2002. The dendritic cell-specific adhesion receptor DC-SIGN internalizes antigen for presentation to T cells. *J. Immunol.* 168: 2118-2126.
2. Geijtenbeek, T.B., et al. 2002. Marginal zone macrophages express a murine homologue of DC-SIGN that captures blood-borne antigens *in vivo*. *Blood* 100: 2908-2916.
3. Moris, A., et al. 2004. DC-SIGN promotes exogenous MHC-I-restricted HIV-1 antigen presentation. *Blood* 103: 2648-2654.
4. Cormier, E.G., et al. 2004. L-SIGN (CD209L) and DC-SIGN (CD209) mediate transinfection of liver cells by hepatitis C virus. *Proc. Natl. Acad. Sci. USA* 101: 14067-14072.
5. Weber, K.S., et al. 2004. Sialylation of ICAM-2 on platelets impairs adhesion of leukocytes via LFA-1 and DC-SIGN. *Inflammation* 28:177-188.
6. Sakuntabhai, A., et al. 2005. A variant in the CD209 promoter is associated with severity of dengue disease. *Nat. Genet.* 37: 507-513.
7. de la Rosa, G., et al. 2005. Regulated recruitment of DC-SIGN to cell-cell contact regions during zymosan-induced human dendritic cell aggregation. *J. Leukoc. Biol.* 77: 699-709.
8. Groot, F., et al. 2005. Lactoferrin prevents dendritic cell-mediated human immunodeficiency virus type 1 transmission by blocking the DC-SIGN—gp120 interaction. *J. Virol.* 79: 3009-3015.

CHROMOSOMAL LOCATION

Genetic locus: Cd209c (mouse) mapping to 8 A1.1.

PRODUCT

CD209c siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CD209c shRNA Plasmid (m): sc-142186-SH and CD209c shRNA (m) Lentiviral Particles: sc-142186-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CD209c siRNA (m) is recommended for the inhibition of CD209c expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CD209c gene expression knockdown using RT-PCR Primer: CD209c (m)-PR: sc-142186-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Jung, T.W., et al. 2018. LECT2 promotes inflammation and Insulin resistance in adipocytes via P38 pathways. *J. Mol. Endocrinol.* 61: 37-45.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.