



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)





Cdc14a phosphatase siRNA (m): sc-142206

BACKGROUND

In budding yeast, the Cdc14a phosphatase activates mitotic exit by dephosphorylation of specific cyclin-dependent kinase (Cdk) substrates and seems to be regulated by sequestration in the nucleolus until its release during mitosis. Human Cdc14a phosphatase is highly similar to *Saccharomyces cerevisiae* Cdc14 and is a member of the dual specificity protein Tyrosine phosphatase family. It interacts with and dephosphorylates tumor suppressor protein p53 and may regulate the function of p53. In addition, Cdc14a dephosphorylates hCdh1 and activates APCdh1. Cdc14a phosphatase plays a role in the regulation of the centrosome cycle, mitosis and cytokinesis, thereby influencing chromosome partitioning and genomic stability in human cells. Deregulated human Cdc14a phosphatase disrupts centrosome separation and chromosome segregation.

REFERENCES

1. Wong, A.K., et al. 1999. Genomic structure, chromosomal location, and mutation analysis of the human CDC14A gene. *Genomics* 59: 248-251.
2. Li, L., et al. 2000. The human Cdc14 phosphatases interact with and dephosphorylate the tumor suppressor protein p53. *J. Biol. Chem.* 275: 2410-2414.
3. Bembek, J., et al. 2001. Regulation of the anaphase-promoting complex by the dual specificity phosphatase human Cdc14a. *J. Biol. Chem.* 276: 48237-48242.
4. Kaiser, B.K., et al. 2002. Disruption of centrosome structure, chromosome segregation, and cytokinesis by misexpression of human Cdc14A Phosphatase. *Mol. Biol. Cell* 13: 2289-2300.
5. Mailand, N., et al. 2002. Deregulated human Cdc14A phosphatase disrupts centrosome separation and chromosome segregation. *Nat. Cell Biol.* 4: 317-322.

CHROMOSOMAL LOCATION

Genetic locus: Cdc14a (mouse) mapping to 3 G1.

PRODUCT

Cdc14a phosphatase siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Cdc14a phosphatase shRNA Plasmid (m): sc-142206-SH and Cdc14a phosphatase shRNA (m) Lentiviral Particles: sc-142206-V as alternate gene silencing products.

For independent verification of Cdc14a phosphatase (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142206A, sc-142206B and sc-142206C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Cdc14a phosphatase siRNA (m) is recommended for the inhibition of Cdc14a phosphatase expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Cdc14a phosphatase gene expression knockdown using RT-PCR Primer: Cdc14a phosphatase (m)-PR: sc-142206-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.