



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# CGI-121 siRNA (m): sc-142305

## BACKGROUND

CGI-121, also known as TP53RK-binding protein and PRPK-binding protein, is a 175 amino acid protein that is widely expressed. Localizing to the cytoplasm and nucleus, CGI-121 interacts with PRPK, a protein kinase that phosphorylates Ser15 of p53. PRPK phosphorylation of p53 causes increased stabilization and activity of p53. CGI-121 may act as an inhibitor of the PRPK-p53 interaction, thus preventing the phosphorylation of p53. The gene encoding CGI-121 maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome. There are three isoforms of CGI-121 that are produced as a result of alternative splicing events.

## REFERENCES

1. Lai, C.H., et al. 2000. Identification of novel human genes evolutionarily conserved in *Caenorhabditis elegans* by comparative proteomics. *Genome Res.* 10: 703-713.
2. Miyoshi, A., et al. 2003. Identification of CGI-121, a novel PRPK (p53-related protein kinase)-binding protein. *Biochem. Biophys. Res. Commun.* 303: 399-405.
3. Facchin, S., et al. 2003. Functional homology between yeast piD261/Bud32 and human PRPK: both phosphorylate p53 and PRPK partially complements piD261/Bud32 deficiency. *FEBS Lett.* 549: 63-66.
4. Abe, Y., et al. 2006. A Small Ras-like protein Ray/Rab1c modulates the p53-regulating activity of PRPK. *Biochem. Biophys. Res. Commun.* 344: 377-385.
5. Wang, X.F., et al. 2006. Identification of differentially expressed genes induced by angiotensin II in rat cardiac fibroblasts. *Clin. Exp. Pharmacol. Physiol.* 33: 41-46.
6. Oppermann, F.S., et al. 2009. Large-scale proteomics analysis of the human kinome. *Mol. Cell Proteomics* 8: 1751-1764.

## CHROMOSOMAL LOCATION

Genetic locus: Tprkb (mouse) mapping to 6 C3.

## PRODUCT

CGI-121 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CGI-121 shRNA Plasmid (m): sc-142305-SH and CGI-121 shRNA (m) Lentiviral Particles: sc-142305-V as alternate gene silencing products.

For independent verification of CGI-121 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142305A, sc-142305B and sc-142305C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

CGI-121 siRNA (m) is recommended for the inhibition of CGI-121 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CGI-121 gene expression knockdown using RT-PCR Primer: CGI-121 (m)-PR: sc-142305-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.