



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Clca6 siRNA (m): sc-142372

BACKGROUND

The calcium-activated chloride channel (CLCA) protein family, which includes the human homologs CLCA1 and CLCA2, display distinct tissue distribution patterns. CLCA1 is expressed as a precursor protein that is processed into two cell surface associated subunits. CLCA1 is upregulated by interleukin-9 and regulates the expression of mucins. CLCA1 may provide a therapeutic target to control mucus overproduction in airway disease patients with cystic fibrosis. CLCA2 expression is downregulated in breast cancer and, therefore, is thought to act as a tumor suppressor in normal cells. CLCA3 is a structurally divergent member of the CLCA family that does not function as a channel protein. CLCA4 is a CLCA member that is expressed in human rectal mucosa, CLCA5 shows strong expression in eye and spleen, and CLCA6 is primarily expressed in intestine and stomach. The mouse homolog, Clca6, is a 925 amino acid protein that is associated with the microvilli of non-goblet cell enterocytes in the murine small and large intestine but in no other tissues. Clca6 may have a direct or indirect role transepithelial anion conductance in the mouse intestine.

REFERENCES

1. Gandhi, R., et al. 1998. Molecular and functional characterization of a calcium-sensitive chloride channel from mouse lung. *J. Biol. Chem.* 273: 32096-32101.
2. Gruber, A.D., et al. 1999. Molecular cloning and transmembrane structure of hCLCA2 from human lung, trachea, and mammary gland. *Am. J. Physiol.* 276: C1261-1270.
3. Gruber, A.D., et al. 1999. Genomic cloning, molecular characterization, and functional analysis of human CLCA1, the first human member of the family of Ca²⁺-activated Cl⁻ channel proteins. *Genomics* 54: 200-214.
4. Hauber, H.P., et al. 2003. Increased expression of interleukin-9, interleukin-9 receptor, chloride channel hCLCA1 in the upper airways of patients with cystic fibrosis. *Laryngoscope* 113: 1037-1042.
5. Li, X., et al. 2004. CLCA2 tumour suppressor gene in 1p31 is epigenic cancer. *Oncogene* 23: 1474-1480.
6. Evans, S.R., et al. 2004. Molecular and functional analyses of two new calcium-activated chloride channel family members from mouse eye and intestine. *J. Biol. Chem.* 279: 41792-41800.
7. Beckley, J.R., et al. 2004. Re-expression of detachment-inducible chloride channel mCLCA5 suppresses growth of metastatic breast cancer cells. *J. Biol. Chem.* 279: 41634-41641.
8. Bothe, M.K., et al. 2008. Murine mCLCA6 is an integral apical membrane protein of non-goblet cell enterocytes and co-localizes with the cystic fibrosis transmembrane conductance regulator. *J. Histochem. Cytochem.* 56: 495-509.
9. Song, J., et al. 2009. Cloning and characterization of a calcium-activated chloride channel in rat uterus. *Biol. Reprod.* 80: 788-794.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Clca6 (mouse) mapping to 3 H2.

PRODUCT

Clca6 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Clca6 shRNA Plasmid (m): sc-142372-SH and Clca6 shRNA (m) Lentiviral Particles: sc-142372-V as alternate gene silencing products.

For independent verification of Clca6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142372A, sc-142372B and sc-142372C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Clca6 siRNA (m) is recommended for the inhibition of Clca6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Clca6 gene expression knockdown using RT-PCR Primer: Clca6 (m)-PR: sc-142372-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.