



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



connexin 31.9 siRNA (m): sc-142495

BACKGROUND

The connexin family of proteins form hexameric complexes, called connexons, that facilitate movement of low molecular weight proteins between cells via gap junctions. Connexin proteins share a common topology of four transmembrane α -helical domains, two extracellular loops, a cytoplasmic loop and cytoplasmic N- and C-termini. Many of the key functional differences arise from specific amino acid substitutions in the most highly conserved domains; the transmembrane and extracellular regions. Connexin 31.9, also known as GJD3 (Gap junction delta-3 protein), is a 294 amino acid protein that interacts with ZO-1, a tight junction protein. Connexin 31.9 is expressed in heart, brain, kidney, spleen, lung, testis, colon and vascular smooth muscle cells. There are two isoforms of connexin 31.9 that are produced as a result of alternative splicing events.

REFERENCES

1. Belluardo, N., White, T.W., Srinivas, M., Trovato-Salinaro, A., Ripps, H., Mudò, G., Bruzzone, R. and Condorelli, D.F. 2001. Identification and functional expression of HCx31.9, a novel gap junction gene. *Cell Commun. Adhes.* 8: 173-178.
2. Nielsen, P.A., Beahm, D.L., Giepmans, B.N., Baruch, A., Hall, J.E. and Kumar, N.M. 2002. Molecular cloning, functional expression, and tissue distribution of a novel human gap junction-forming protein, connexin-31.9. Interaction with zona occludens protein-1. *J. Biol. Chem.* 277: 38272-38283.
3. Nielsen, P.A. and Kumar, N.M. 2003. Differences in expression patterns between mouse connexin-30.2 (Cx30.2) and its putative human orthologue, connexin-31.9. *FEBS Lett.* 540: 151-156.
4. Miquerol, L., Dupays, L., Theveniau-Ruissy, M., Alcolea, S., Jarry-Guichard, T., Abran, P. and Gros, D. 2003. Gap junctional connexins in the developing mouse cardiac conduction system. *Novartis Found. Symp.* 250: 80-98.
5. Cruciani, V. and Mikalsen, S.O. 2005. The connexin gene family in mammals. *Biol. Chem.* 386: 325-332.
6. Bukauskas, F.F., Kreuzberg, M.M., Rackauskas, M., Bukauskiene, A., Bennett, M.V., Verselis, V.K. and Willecke, K. 2006. Properties of mouse connexin 30.2 and human connexin 31.9 hemichannels: implications for atrioventricular conduction in the heart. *Proc. Natl. Acad. Sci. USA* 103: 9726-9731.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 607425. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Renner, R., Paasch, U., Simon, J.C., Froster, U.G. and Heinritz, W. 2008. A new mutation in the GJB3 gene in a patient with erythrokeratoderma variabilis. *J. Eur. Acad. Dermatol. Venereol.* 22: 750-751.
9. Herve, J.C., Derangeon, M., Theveniau-Ruissy, M., Miquerol, L., Sarrouilhe, D. and Gros, D. 2008. Connexins and junctional channels. Roles in the spreading of cardiac electrical excitation and heart development. *Pathol. Biol.* 56: 334-341.

CHROMOSOMAL LOCATION

Genetic locus: Gjd3 (mouse) mapping to 11 D.

PRODUCT

connexin 31.9 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see connexin 31.9 shRNA Plasmid (m): sc-142495-SH and connexin 31.9 shRNA (m) Lentiviral Particles: sc-142495-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

connexin 31.9 siRNA (m) is recommended for the inhibition of connexin 31.9 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor connexin 31.9 gene expression knockdown using RT-PCR Primer: connexin 31.9 (m)-PR: sc-142495-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.