



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CRBP II siRNA (m): sc-142563

BACKGROUND

The cellular retinol-binding proteins (CRBP I, II, III and IV) belong to a super-family of small cytoplasmic proteins that interact with hydrophobic ligands. Vitamin A, a molecule essential for cell growth and differentiation, embryonic development and vision, is transported into the cell by the CRBPs in its alcoholic form, called retinol. Both CRBP I and II are composed of ten antiparallel β -strands, which form a β -barrel that contains the retinol molecule, and two α -helices, which cover the open ends of the barrel. CRBP II, which is also known as RBP2 (retinol-binding protein 2), consists of 134 amino acids and is expressed solely in the small intestine where it mediates the absorption of retinoids and carotenoids to biosynthesize retinyl esters.

REFERENCES

1. Ong, D.E. and Page, D.L. 1986. Quantitation of cellular retinol-binding protein in human organs. *Am. J. Clin. Nutr.* 44: 425-430.
2. Cowan, S.W., Newcomer, M.E. and Jones, T.A. 1993. Crystallographic studies on a family of cellular lipophilic transport proteins. Refinement of P2 myelin protein and the structure determination and refinement of cellular retinol-binding protein in complex with all-trans-retinol. *J. Mol. Biol.* 230: 1225-1246.
3. Winter, N.S., Bratt, J.M. and Banaszak, L.J. 1993. Crystal structures of holo- and apo-cellular retinol-binding protein II. *J. Mol. Biol.* 230: 1247-1259.
4. Okuno, M., Numaguchi, S., Moriwaki, H. and Muto, Y. 1993. Cellular retinoid-binding proteins. *Nippon Rinsho* 51: 879-885.
5. Takase, S., Suruga, K. and Goda, T. 2000. Regulation of vitamin A metabolism-related gene expression. *Br. J. Nutr.* 84: S217-S221.
6. Xu, G., Bochaton-Piallat, M.L., Andreutti, D., Low, R.B., Gabbiani, G. and Neuville, P. 2001. Regulation of α -smooth muscle Actin and CRBP-1 expression by retinoic acid and TGF β in cultured fibroblasts. *J. Cell. Physiol.* 187: 315-325.
7. Folli, C., Calderone, V., Ottonello, S., Bolchi, A., Zanotti, G., Stoppini, M. and Berni, R. 2001. Identification, retinoid binding and x-ray analysis of a human retinol-binding protein. *Proc. Natl. Acad. Sci. USA* 98: 3710-3715.

CHROMOSOMAL LOCATION

Genetic locus: Rbp2 (mouse) mapping to 9 E3.3.

PRODUCT

CRBP II siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CRBP II shRNA Plasmid (m): sc-142563-SH and CRBP II shRNA (m) Lentiviral Particles: sc-142563-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CRBP II siRNA (m) is recommended for the inhibition of CRBP II expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CRBP II gene expression knockdown using RT-PCR Primer: CRBP II (m)-PR: sc-142563-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.