



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CYB5D2 siRNA (m): sc-142651

BACKGROUND

Cytochrome b5 is a membrane-bound heme protein that possesses two heme groups and functions as an electron carrier for many membrane-bound oxygenases. Two isoforms of cytochrome b5, a microsomal membrane-bound form and a cytoplasmic form, are produced by alternative splicing. Mutations in the gene encoding cytochrome b5 are associated with Leber's hereditary optic neuropathy and myopathy. CYB5D2 (cytochrome b5 domain containing 2) is a 264 amino acid single-pass membrane protein that contains one cytochrome b5 heme-binding domain, suggesting a similar function to cytochrome b5. The gene encoding CYB5D2 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

REFERENCES

1. Abe, K., Kimura, S., Kizawa, R., Anan, F.K. and Sugita, Y. 1985. Amino acid sequences of cytochrome b5 from human, porcine, and bovine erythrocytes and comparison with liver microsomal cytochrome b5. *J. Biochem.* 97: 1659-1668.
2. Yoo, M. and Steggles, A.W. 1988. The complete nucleotide sequence of human liver cytochrome b5 mRNA. *Biochem. Biophys. Res. Commun.* 156: 576-580.
3. Giordano, S.J. and Steggles, A.W. 1991. The human liver and reticulocyte cytochrome b5 mRNAs are products from a single gene. *Biochem. Biophys. Res. Commun.* 178: 38-44.
4. Hom, K., Ma, Q.F., Wolfe, G., Zhang, H., Storch, E.M., Daggett, V., Basus, V.J. and Waskell, L. 2000. NMR studies of the association of cytochrome b5 with cytochrome c. *Biochemistry* 39: 14025-14039.
5. Wang, Y.H., Ren, Y., Wang, W.H., Xie, Y. and Huang, Z.X. 2001. The regulation of surface charged residues on the properties of cytochrome b5. *J. Protein Chem.* 20: 487-493.
6. Mapes, S., Tarantal, A.F., Parker, C.R., Moran, F.M., Bahr, J.M., Pyter, L. and Conley, A.J. 2002. Adrenocortical cytochrome b5 expression during fetal development of the rhesus macaque. *Endocrinology* 143: 1451-1458.
7. Reed, J.R. and Hollenberg, P.F. 2003. Examining the mechanism of stimulation of cytochrome P450 by cytochrome b5: the effect of cytochrome b5 on the interaction between cytochrome P450 2B4 and P450 reductase. *J. Inorg. Biochem.* 97: 265-275.
8. Zhang, Q., Cao, C., Wang, Z.Q., Wang, Y.H., Wu, H. and Huang, Z.X. 2004. The comparative study on the solution structures of the oxidized bovine microsomal cytochrome b5 and mutant V45H. *Protein Sci.* 13: 2161-2169.
9. Sun, N., Wang, A., Cowley, A.B., Altuve, A., Rivera, M. and Benson, D.R. 2005. Enhancing the stability of microsomal cytochrome b5: a rational approach informed by comparative studies with the outer mitochondrial membrane isoform. *Protein Eng. Des. Sel.* 18: 571-579.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Cyb5d2 (mouse) mapping to 11 B4.

PRODUCT

CYB5D2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CYB5D2 shRNA Plasmid (m): sc-142651-SH and CYB5D2 shRNA (m) Lentiviral Particles: sc-142651-V as alternate gene silencing products.

For independent verification of CYB5D2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142651A, sc-142651B and sc-142651C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CYB5D2 siRNA (m) is recommended for the inhibition of CYB5D2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CYB5D2 gene expression knockdown using RT-PCR Primer: CYB5D2 (m)-PR: sc-142651-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.