



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cyp3a16 siRNA (m): sc-142713

BACKGROUND

The cytochrome P450 proteins (Cyps) are monooxygenases that catalyze reactions involved in both drug metabolism and in the synthesis of cholesterol, steroids and other lipids. P450 enzymes are classified into subfamilies, including Cyp1, Cyp2 and Cyp3, based on their sequence similarities. In mice, there are ten Cyp3 genes which encode proteins that are designated Cyp3a11, Cyp3a13, Cyp3a16, Cyp3a25, Cyp3a91, Cyp3a41a, Cyp3a41b, Cyp3a44, Cyp3a57 and Cyp3a59. Most of the mouse Cyp3 proteins are localized to the endoplasmic reticulum and function to catalyze reactions related to the synthesis, degradation and detoxification of various compounds.

REFERENCES

1. Yanagimoto, T., et al. 1992. Mouse liver cytochrome P-450 (P-450IIAM1): its cDNA cloning and inducibility by dexamethasone. *Biochim. Biophys. Acta* 1130: 329-332.
2. Ramana, K.V., et al. 1998. Gene regulation of cytochrome P450—an overview. *Indian J. Exp. Biol.* 36: 437-446.
3. Nallani, S.C., et al. 2001. Increased activity of CYP3A enzyme in primary cultures of rat hepatocytes treated with docetaxel: comparative evaluation with paclitaxel. *Cancer Chemother. Pharmacol.* 48: 115-122.
4. McArthur, A.G., et al. 2003. Phylogenetic analysis of the cytochrome P450 3 (CYP3) gene family. *J. Mol. Evol.* 57: 200-211.
5. Lewis, D.F., et al. 2004. Compound lipophilicity for substrate binding to human P450s in drug metabolism. *Drug Discov. Today* 9: 530-537.
6. Murray, M., et al. 2004. Comparative inhibition of inducible and constitutive CYPs in rat hepatic microsomes by parathion. *Xenobiotica* 34: 723-739.
7. Girault, I., et al. 2005. Simultaneous measurement of 23 isoforms from the human cytochrome P450 families 1 to 3 by quantitative reverse transcriptase-polymerase chain reaction. *Drug Metab. Dispos.* 33: 1803-1810.
8. Ioannides, C. 2006. Cytochrome p450 expression in the liver of food-producing animals. *Curr. Drug Metab.* 7: 335-348.

CHROMOSOMAL LOCATION

Genetic locus: Cyp3a16 (mouse) mapping to 5 G2.

PRODUCT

Cyp3a16 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Cyp3a16 shRNA Plasmid (m): sc-142713-SH and Cyp3a16 shRNA (m) Lentiviral Particles: sc-142713-V as alternate gene silencing products.

For independent verification of Cyp3a16 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142713A, sc-142713B and sc-142713C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Cyp3a16 siRNA (m) is recommended for the inhibition of Cyp3a16 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Cyp3a16 gene expression knockdown using RT-PCR Primer: Cyp3a16 (m)-PR: sc-142713-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.