



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Cyp4f17 siRNA (m): sc-142733

BACKGROUND

Cytochrome P450 proteins are heme-thiolate monooxygenases that mediate NADPH-dependent electron transport and function to oxidize a variety of structurally unrelated compounds, including steroids, fatty acids and xenobiotics. Specifically, cytochrome P450s are responsible for metabolizing arachidonic acid to hydroxyeicosatetraenoic acid (a regulator of blood pressure) and epoxyeicosatrienoic acid (a molecule involved in signaling events). Cyp4f17 (cytochrome P450, family 4, subfamily f, polypeptide 17), also known as EG208285, is a 524 amino acid protein that is encoded by a gene mapping to mouse chromosome 17 B1.

REFERENCES

1. Simpson, A.E. 1997. The cytochrome P450 4 (CYP4) family. *Gen. Pharmacol.* 28: 351-359.
2. Bylund, J., Finnström, N. and Oliw, E.H. 1999. Gene expression of a novel cytochrome P450 of the CYP4F subfamily in human seminal vesicles. *Biochem. Biophys. Res. Commun.* 261: 169-174.
3. Bylund, J., Hidestrand, M., Ingelman-Sundberg, M. and Oliw, E.H. 2000. Identification of CYP4F8 in human seminal vesicles as a prominent 19-hydroxylase of prostaglandin endoperoxides. *J. Biol. Chem.* 275: 21844-21849.
4. Oliw, E.H., Stark, K. and Bylund, J. 2001. Oxidation of prostaglandin H₂ and prostaglandin H₂ analogues by human cytochromes P450: analysis of ω -side chain hydroxy metabolites and four stereoisomers of 5-hydroxyprostaglandin I₁ by mass spectrometry. *Biochem. Pharmacol.* 62: 407-415.
5. Stark, K., Törmä, H., Cristea, M. and Oliw, E.H. 2003. Expression of CYP4F8 (prostaglandin H 19-hydroxylase) in human epithelia and prominent induction in epidermis of psoriatic lesions. *Arch. Biochem. Biophys.* 409: 188-196.
6. Nelson, D.R., Zeldin, D.C., Hoffman, S.M., Maltais, L.J., Wain, H.M. and Nebert, D.W. 2004. Comparison of cytochrome P450 (CYP) genes from the mouse and human genomes, including nomenclature recommendations for genes, pseudogenes and alternative-splice variants. *Pharmacogenetics* 14: 1-18.
7. Stark, K., Wongsud, B., Burman, R. and Oliw, E.H. 2005. Oxygenation of polyunsaturated long chain fatty acids by recombinant CYP4F8 and CYP4F12 and catalytic importance of Tyr-125 and Gly-328 of CYP4F8. *Arch. Biochem. Biophys.* 441: 174-181.
8. Stark, K., Bylund, J., Törmä, H., Sahlen, G. and Oliw, E.H. 2005. On the mechanism of biosynthesis of 19-hydroxyprostaglandins of human seminal fluid and expression of cyclooxygenase-2, PGH 19-hydroxylase (CYP4F8) and microsomal PGE synthase-1 in seminal vesicles and vas deferens. *Prostaglandins Other Lipid Mediat.* 75: 47-64.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Cyp4f17 (mouse) mapping to 17 B1.

PRODUCT

Cyp4f17 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Cyp4f17 shRNA Plasmid (m): sc-142733-SH and Cyp4f17 shRNA (m) Lentiviral Particles: sc-142733-V as alternate gene silencing products.

For independent verification of Cyp4f17 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-142733A and sc-142733B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Cyp4f17 siRNA (m) is recommended for the inhibition of Cyp4f17 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Cyp4f17 gene expression knockdown using RT-PCR Primer: Cyp4f17 (m)-PR: sc-142733-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.